arXiv:1206.2531v1 [math.RT] 12 Jun 2012

A GABRIEL-TYPE THEOREM FOR CLUSTER TILTING

YUYA MIZUNO

ABSTRACT. We study the relationship between n-cluster tilting modules over n representation finite algebras and the Euler forms. We show that the dimension vectors of cluster-indecomposable modules give the roots of the Euler form. Moreover, we show that cluster-indecomposable modules are uniquely determined by their dimension vectors. This is a generalization of Gabriel's theorem by cluster tilting theory. We call the above roots cluster-roots and investigate their properties. Furthermore, we provide the description of quivers with relations of n-APR tilts. Using this, we provide a generalization of BGP reflection functors.

CONTENTS

1. Introduction 1
2. Preliminaries 4
2.1. Euler forms, Cartan matrix and Coxeter transformation 4
2.2. n-representation-finite algebras 5
2.3. n-APR tilting modules 6
3. Cluster-indecomposable modules and their properties 8
3.1. Cluster-roots 8
3.2. Reflection transformations 10
4. Cluster-roots determine cluster-indecomposable modules 13
5. A criterion for cluster-roots 19
6. Quivers of n-APR tilts and reflection functors 24
References 29

1. INTRODUCTION

Global dimension is one of the important invariants of algebras and it allows to measure the complexity of algebras. The algebras with global dimension 0 are semisimple. The next fundamental class consists of the algebras with global dimension at most 1, which are characterized as path algebras up to Morita equivalence. With regard to this class, Gabriel gave the following famous theorem in 1972.

Theorem 1.1 (Gabriel). *Let K be an algebraically closed field, Q be a connected Dynkin quiver and q_Q be the Euler form (Definition 2.1) of the path algebra KQ.*

(1) *For any indecomposable KQ-module X, the dimension vector $\underline{\dim}X$ is a positive root of q_Q.*
(2) *The map $X \mapsto \underline{\dim}X$ gives a bijection between the isomorphism classes of indecomposable KQ-modules and the positive roots of q_Q.*

The author is supported by Grant-in-Aid for JSPS Fellowships No.23.5593.

A generalization of this notable theorem was given by many authors. One of them was given by [DF, N, DR, Ka], where path algebras of non-Dynkin quivers were studied, and another was given by [B2, D], where representation-directed algebras with global dimension 2 were studied. Let us point out that the notion of reflection functors [BGP] was inspired by the theorem and it led to tilting theory [APR, BB, HR1, M, H, Ric], which is an important tool in modern representation theory of algebras. Thus this theorem plays a significant role in representation theory of algebras.

The aim of this paper is to generalize Gabriel's theorem from the viewpoint of higher dimensional Auslander-Reiten theory, which was introduced by Iyama [I1, I2, I3] and has been developed by several authors [EH, HI1, HI2, HIO, HZ, IO1, IO2]. It is also closely related to the recent studies of cluster tilting theory (see for example [Ke1, Ke2, Re]). In higher dimensional Auslander-Reiten theory, *n-representation-finite algebras* (Definition 2.3) are fundamental objects. Recall that we call an algebra n-representation-finite if its global dimension is at most n and it has an n-cluster tilting module, and we call a Λ-module *cluster-indecomposable* if it is isomorphic to an indecomposable direct summand of an n-cluster tilting module. Note that 1-representation-finite algebras are exactly path algebras of Dynkin quivers and any indecomposable module is cluster-indecomposable in this case (Remark 2.4).

Let us investigate path algebras over a Dynkin quiver in the context of Auslander-Reiten theory. For a Dynkin quiver Q, we have the following, where τ and τ^- are Auslander-Reiten translations.

- Any indecomposable KQ-module X is isomorphic to $\tau^i I$ for an integer $i \geq 0$ and indecomposable injective KQ-module I.
- Any indecomposable KQ-module X is isomorphic to $\tau^{-i} P$ for an integer $i \geq 0$ and indecomposable projective KQ-module P.

Then for an n-representation-finite algebra Λ, we have the following (Proposition 2.5), where τ_n and τ_n^- are n-Auslander-Reiten translations.

- Any cluster-indecomposable Λ-module X is isomorphic to $\tau_n^i I$ for an integer $i \geq 0$ and indecomposable injective Λ-module I.
- Any cluster-indecomposable Λ-module X is isomorphic to $\tau_n^{-i} P$ for an integer $i \geq 0$ and indecomposable projective Λ-module P.

We focus on this property and give a Gabriel-type theorem for algebras with the global dimension n.

Our main results of the paper are given by the following theorem.

Theorem 1.2 (Theorem 3.2, 4.12)**.** *Let Λ be an n-representation-finite algebra and q_Λ be the Euler form of Λ.*

(1) *For any cluster-indecomposable Λ-module X, the dimension vector $\underline{\dim} X$ is a positive root of q_Λ.*
(2) *The map $X \mapsto \underline{\dim} X$ gives a injection between the isomorphism classes of cluster-indecomposable Λ-modules and the positive roots of q_Λ.*

We call the above positive roots *cluster-roots* (Definition 3.3). Thus cluster-roots correspond bijectively to cluster-indecomposable modules, which is an analogue of Theorem 1.1. In fact, when $n = 1$, cluster-roots and cluster-indecomposable modules coincide with positive roots and indecomposable modules, respectively. On the other hand, the map is not necessarily surjective if $n > 1$. Therefore our next aim is to characterize cluster-roots.

We give a complete description of cluster-roots by *reflection transformations* (Definition 3.6), which are linear transformations of the Grothendieck group.

Theorem 1.3 (Theorem 4.10)**.** *Let* Λ *be an* n*-representation-finite algebra and* Q *be the quiver of* Λ*. Assume that* Q *has an admissible ordering* $(1, \dots, l)$ *(Definition 2.12). We denote by* t_i *the reflection transformation associated with vertex* i *and* $\mathbf{c} := t_l \dots t_1$*.*

1. *Any cluster-root can be written as* $\mathbf{c}^{-u}\mathbf{p}_i$ *for an integer* $u \geq 0$ *and* $i \in Q_0$*, where* $\mathbf{p}_i := t_1^{-1} \dots t_{i-1}^{-1}\mathbf{e}_i$*.*
2. *Any cluster-root can be written as* $\mathbf{c}^{v}\mathbf{q}_j$ *for an integer* $v \geq 0$ *and* $j \in Q_0$*, where* $\mathbf{q}_j := t_l \dots t_{j+1}\mathbf{e}_j$*.*

Thus cluster-roots can be calculated in terms of linear algebra. Furthermore, we give the following terminology to characterize cluster-roots.

Definition 1.4 (Definition 5.1)**.** We call $x \in \mathbb{Z}^l$ Φ-*sign-coherent* if $\Phi^m(x) \in \mathbb{Z}^l_{\geq 0}$ or $\Phi^m(x) \in \mathbb{Z}^l_{\leq 0}$ for any $m \in \mathbb{Z}$ and call $x \in \mathbb{Z}^l$ Φ-*positive* if $\Phi^m(x) \in \mathbb{Z}^l_{\geq 0}$ for any $m \in \mathbb{Z}$, where Φ is the Coxeter transformation (Definition 2.1).

Then we have the following result.

Theorem 1.5 (Theorem 5.2)**.** *Cluster-roots are* Φ*-sign-coherent. Moreover, if* n *is even, then cluster-roots are* Φ*-positive.*

We consider the converse for $n = 2$ and pose the following conjecture.

Conjecture 1.6 (Conjecture 5.3)**.** *Let* Λ *be a* 2*-representation-finite algebra. Then* Φ*-positive roots are cluster-roots. This is equivalent, by Theorem 1.2, to that* Φ*-positive roots correspond bijectively to the isomorphism classes of cluster-indecomposable modules.*

We prove the conjecture for some classes of algebras.

Theorem 1.7 (Corollary 5.8)**.** *If* Λ *is iterated tilted, then Conjecture 1.6 is true.*

We now describe the organization of this paper.

In section 2, we recall basic definitions and results about n-representation-finite algebras and n-APR tilting modules. An n-APR tilting module over an n-representation-finite algebra induces an equivalence between certain subcategories (Corollary 2.11). This property plays a key role in this paper.

In section 3.1, we explain the connection between the Coxeter transformation and n-AR translations. In particular, we show that the dimension vectors of cluster-indecomposable modules are roots of the Euler form. In section 3.2, we introduce reflection transformations, which are linear transformations of the Grothendieck group. We will show that, in a nice subcategory, reflection transformations are compatible with the reflection functors.

In section 4, we define the Coxeter functors as a composition of reflection functors. Using them, we show that cluster-indecomposable modules can be constructed from a simple module by applying a sequence of the reflection functors. This is a generalization of well-known properties of BGP reflection functors. As a corollary, we give a description of cluster-roots in terms of reflection transformations. Moreover we prove that cluster-indecomposable modules are uniquely determined by their dimension vectors.

In section 5, we study a characterization of cluster-roots. We introduce the notion of Φ-sign-coherence and Φ-positive. We show that cluster-roots are Φ-sign-coherent roots. Moreover, we conjecture that Φ-positive roots are cluster-roots for $n = 2$. We show that this is true for a certain class of algebras.

Section 6 is independent from the previous sections. We give descriptions of n-APR tilts in terms of quivers with relations. Moreover, following [BGP], we give an interpretation of reflection functors in terms of linear representations of quivers. Then we explain the connection with the functors defined by n-APR tilting modules.

Notations Let K be an algebraically closed field and we denote by $D := \mathrm{Hom}_K(-, K)$. By an algebra Λ, we mean a basic, connected, finite dimensional algebra over K. All modules are right modules. We denote by $\mathrm{mod}\,\Lambda$ the category of finitely generated Λ-modules and by $\mathrm{add}M$ the subcategory of $\mathrm{mod}\,\Lambda$ consisting of direct summands of finite direct sums of copies of M. The composition gf means first f, then g. We always assume that a quiver Q is finite, connected, acyclic with the number of vertices l (denote by $|Q_0| = l$). We denote by Q_0 vertices and Q_1 arrows of Q and by $a : s(a) \to e(a)$ the start and end vertices of an arrow or path a. We denote by $KQ/(R)$ a path algebra KQ bound by a set of relations R. We denote by P_i, I_i, S_i, respectively, the indecomposable projective, injective, simple $KQ/(R)$-module corresponding to the vertex $i \in Q_0$.

Acknowledgement First and foremost, the author would like to thank Osamu Iyama for his support and patient guidance. Part of this work was done while he visited Trondheim. He would like to thank Idun Reiten and Steffen Oppermann for their suggestions and kind advice for correcting his English, and thank all the people at NTNU for their hospitality. He is also grateful to Martin Herschend, Kota Yamaura and Takahide Adachi for their valuable comments and advice.

2. Preliminaries

In this section, we give a summary of definitions and results we will use in this paper. Throughout this paper, we always assume that the quiver Q is finite, connected, acyclic.

2.1. **Euler forms, Cartan matrix and Coxeter transformation.** We recall some terminology. We refer to [ASS, Rin1] for details.

Definition 2.1. Let Λ be a basic finite dimensional algebra with a complete set $\{e_1, \ldots, e_l\}$ of primitive orthogonal idempotents. Assume that Λ has finite global dimension n.

- The *Cartan matrix* of Λ is the (l, l)-matrix

$$C_\Lambda = (\dim(e_i \Lambda e_j))_{1 \le i,j \le l}.$$

The *Coxeter transformation* of Λ is defined to be the matrix

$$\Phi_\Lambda = \Phi := (-1)^n C_\Lambda^t C_\Lambda^{-1}.$$

Note that this definition is slightly different from the ordinary Coxeter transformation (cf. [ASS, 3.14. Definition]). The difference is crucial in Lemma 3.1.

- The *Euler form* of Λ is the $\mathbb{Z}$-bilinear form

$$\langle -, - \rangle : \mathbb{Z}^l \times \mathbb{Z}^l \to \mathbb{Z}$$

defined by $\langle x, y \rangle := x^t (C_\Lambda^{-1})^t y$ for $x, y \in \mathbb{Z}^l$. We define a quadratic form $q_\Lambda(x) := \langle x, x \rangle$ and call a vector $x \in \mathbb{Z}^l$ *root* if $q_\Lambda(x) = 1$.

- For a Λ-module X, we denote the *dimension vector* by

$$\underline{\dim} X := (\dim(Xe_1), \ldots, \dim(Xe_l)).$$

Then we recall the following properties.

Lemma 2.2. *Let $\Lambda = KQ/(R)$ be a finite dimensional algebra with finite global dimension n.*

1. *We have $(C_\Lambda^t C_\Lambda^{-1})\underline{\dim} P_i = \underline{\dim} I_i$ for any $i \in Q_0$.*
2. *We have $\langle x, y\rangle = (-1)^n \langle y, \Phi(x)\rangle = \langle \Phi(x), \Phi(y)\rangle$.*
3. *For any X, Y in $\operatorname{mod}\Lambda$, we have*

$$\langle \underline{\dim} X, \underline{\dim} Y\rangle = \sum_{j\geq 0} (-1)^j \dim \operatorname{Ext}^j_\Lambda(X,Y).$$

2.2. **n-representation-finite algebras.** In this paper, we mainly deal with n-representation-finite algebras. They were investigated in [HI1, I3, IO1, IO2] and, for $n = 2$, in [HI2]. See also related studies [I1, I2].

Definition 2.3. We call a finite dimensional algebra Λ *n-representation-finite* (for a positive integer n) if $\operatorname{gl.dim}\Lambda \leq n$ and there exists an *n-cluster tilting Λ-module* M, i.e.

$$\begin{aligned} \operatorname{add} M &= \{X \in \operatorname{mod}\Lambda \mid \operatorname{Ext}^i_\Lambda(M,X) = 0 \ \text{ for any } \ 0 < i < n\}, \\ &= \{X \in \operatorname{mod}\Lambda \mid \operatorname{Ext}^i_\Lambda(X,M) = 0 \ \text{ for any } \ 0 < i < n\}. \end{aligned}$$

We call a Λ-module X *cluster-indecomposable* if X is isomorphic to an indecomposable direct summand of M.

Remark 2.4. A basic 1-representation-finite algebra Λ is a path algebra of a Dynkin quiver. Indeed, an algebra Λ is 1-representation-finite if and only if it is representation-finite and hereditary. Then, by Gabriel's Theorem, we have $\Lambda = KQ$ for a Dynkin quiver Q.

Let Λ be a finite dimensional algebra with finite global dimension n. We denote by $\mathcal{D}^b(\operatorname{mod}\Lambda)$ the bounded derived category of $\operatorname{mod}\Lambda$. Then we define the Nakayama functor

$$\nu := D \circ \mathbb{R}\operatorname{Hom}_\Lambda(-,\Lambda) : \mathcal{D}^b(\operatorname{mod}\Lambda) \to \mathcal{D}^b(\operatorname{mod}\Lambda),$$

and we write

$$\nu_n := \nu \circ [-n] : \mathcal{D}^b(\operatorname{mod}\Lambda) \to \mathcal{D}^b(\operatorname{mod}\Lambda).$$

Note that Φ gives the action of ν_n on the Grothendieck group $K_0(\mathcal{D}^b(\operatorname{mod}\Lambda))$. Moreover we define *$n$-Auslander-Reiten translations* by the following functors

$$\begin{aligned} \tau_n &:= D\operatorname{Ext}^n_\Lambda(-,\Lambda) : \operatorname{mod}\Lambda \to \operatorname{mod}\Lambda, \\ \tau_n^- &:= \operatorname{Ext}^n_\Lambda(D\Lambda,-) : \operatorname{mod}\Lambda \to \operatorname{mod}\Lambda. \end{aligned}$$

They are related to the functors ν_n and ν_n^- as follows.

$$\begin{aligned} \tau_n &= H^0(\nu_n -) : \operatorname{mod}\Lambda \to \operatorname{mod}\Lambda, \\ \tau_n^- &= H^0(\nu_n^- -) : \operatorname{mod}\Lambda \to \operatorname{mod}\Lambda. \end{aligned}$$

Then we recall some important properties of n-representation-finite algebras.

Proposition 2.5. [I1, I3] *Let $\Lambda = KQ/(R)$ be an n-representation-finite algebra and M be a basic n-cluster tilting module.*

1. *There exist non-negative integers m_i and a bijection $\sigma : Q_0 \to Q_0$ such that $\tau_n^{m_i} I_i \cong P_{\sigma(i)}$ for any $i \in Q_0$. Moreover, we have $M \cong \bigoplus_{i\geq 0} \tau_n^{-i}(\Lambda) \cong \bigoplus_{i\geq 0} \tau_n^{i}(D\Lambda)$.*
2. *We have an equivalence $\tau_n : \operatorname{add}(M/\Lambda) \to \operatorname{add}(M/D\Lambda)$ with a quasi-inverse $\tau_n^- : \operatorname{add}(M/D\Lambda) \to \operatorname{add}(M/\Lambda)$.*

(3) *There exist the following functorial isomorphisms for any* $X, Y \in \operatorname{add} M$ *and* $Z \in \operatorname{mod} \Lambda$.

$$\underline{\operatorname{Hom}}_\Lambda(X, Z) \cong D \operatorname{Ext}^n_\Lambda(Z, \tau_n X), \quad \overline{\operatorname{Hom}}_\Lambda(Z, Y) \cong D \operatorname{Ext}^n_\Lambda(\tau_n^- Y, Z).$$

(4) *We have* $\operatorname{Hom}_\Lambda(\tau_n^i(D\Lambda), \tau_n^j(D\Lambda)) = 0$ *for any* $i < j$.

(5) τ_n *and* τ_n^- *are nilpotent, i.e.* $\tau_n^N = 0$ *and* $\tau_n^{-N} = 0$ *for a sufficiently large integer* N.

(6) *We have* $\tau_n \cong \nu_n$ *on* $\operatorname{add}(M/\Lambda)$ *and* $\tau_n^- \cong \nu_n^-$ *on* $\operatorname{add}(M/D\Lambda)$.

2.3. **n-APR tilting modules.** BGP reflection functors enable us to give a conceptual proof of Gabriel's theorem [BGP]. These functors are now formulated in terms of APR tilting modules [APR]. A relationship between path algebras of Dynkin quivers and APR tilting modules were generalized in [IO1], where the authors introduced the notion of n-representation-finite algebras and n-APR tilting modules. We briefly recall some definitions and results.

Definition 2.6. [IO1, Definition 3.1] Let Λ be a basic finite dimensional algebra, k be a sink of the quiver of Λ and P_k be a simple projective Λ-module. If $\text{inj.dim} P_k = n$ and $\operatorname{Ext}^i_\Lambda(D\Lambda, P_k) = 0$ for any $0 \leq i < n$, we call

$$T_k := \tau_n^- P_k \oplus \Lambda/P_k$$

an *n-APR tilting module*. Dually we define *n-APR cotilting modules*.

Note that we have an ordinary APR tilting module if $n = 1$.

Recall that we call a Λ-module T *tilting module of projective dimension* n [M] if $\text{proj.dim} T \leq n$, $\operatorname{Ext}^i_\Lambda(T, T) = 0$ for $1 \leq i \leq n$ and there exists an exact sequence $0 \to \Lambda \to T_0 \to \cdots \to T_n \to 0$ with T_i in $\operatorname{add} T$. Then we have the following result.

Proposition 2.7. [IO1, Theorem 3.2] *An n-APR tilting module is a tilting module of projective dimension n.*

It is a classical result that 1-representation-finite algebras (= path algebras of a Dynkin quiver) are closed under taking endomorphism algebras of APR tilting modules. The following proposition is a generalization of this property and shows that n-APR tilting modules behave nicely over n-representation-finite algebras.

Proposition 2.8. [IO1, Theorem 4.2, 4.7] *Let Λ be a basic n-representation-finite algebra and k be a sink of the quiver of Λ.*

(1) P_k *satisfies Definition 2.6 and admits the n-APR tilting modules* T_k.

(2) $\operatorname{End}_\Lambda(T_k)$ *is an n-representation-finite algebra.*

(3) $\operatorname{Ext}^n_\Lambda(T_k, P_k)$ *is isomorphic to the simple injective* $\operatorname{End}_\Lambda(T_k)$*-module corresponding to the vertex* k.

Thus a sink vertex k becomes a source vertex of $\operatorname{End}_\Lambda(T_k)$. A more detailed description of the quiver and relations of $\operatorname{End}_\Lambda(T_k)$ will be given in section 6. By the above proposition, it is natural to ask a relationship between n-cluster tilting modules of $\operatorname{mod} \Lambda$ and $\operatorname{mod} \operatorname{End}_\Lambda(T_k)$. For this purpose, we recall a result of tilting theory.

Let T be a tilting Λ-module and $\Gamma := \operatorname{End}_\Lambda(T)$. Let $\mathcal{D}^b(\operatorname{mod} \Lambda)$ and $\mathcal{D}^b(\operatorname{mod} \Gamma)$ be the bounded derived category of $\operatorname{mod} \Lambda$ and $\operatorname{mod} \Gamma$, respectively. Then we have functors

$$\mathbf{F} := \mathbb{R}\operatorname{Hom}_\Lambda(T, -) : \mathcal{D}^b(\operatorname{mod} \Lambda) \to \mathcal{D}^b(\operatorname{mod} \Gamma),$$

$$\mathbf{F}_i := \operatorname{Ext}^i_\Lambda(T, -) : \operatorname{mod} \Lambda \to \operatorname{mod} \Gamma,$$

and we write

$$\begin{aligned} \mathcal{F}_i(T) = \mathcal{F}_i := & \quad \{X \in \operatorname{mod}\Lambda \mid \operatorname{Ext}^j_\Lambda(T,X) = 0 \text{ for any } j \neq i\}, \\ \mathcal{X}_i(T) = \mathcal{X}_i := & \quad \{Y \in \operatorname{mod}\Gamma \mid \operatorname{Tor}^\Gamma_j(Y,T) = 0 \text{ for any } j \neq i\}. \end{aligned}$$

First we recall the following result of Happel.

Lemma 2.9. [H] *In the above set-up,* $\mathbf{F} : \mathcal{D}^b(\operatorname{mod}\Lambda) \to \mathcal{D}^b(\operatorname{mod}\Gamma)$ *gives an equivalence and the restriction of the functor* $[i] \circ \mathbf{F}$ *gives an equivalence* $\mathbf{F}_i : \mathcal{F}_i \to \mathcal{X}_i$.

Next we give a relationship between the cluster tilting subcategories and $\mathcal{F}_0$ as follows.

Lemma 2.10. *Let* $\Lambda = KQ/(R)$ *be an* n*-representation-finite algebra,* $k \in Q_0$ *be a sink,* T_k *be the corresponding* n*-APR tilting* Λ*-module and* X *be a cluster-indecomposable* Λ*-module. Then exactly one of the following holds.*

(1) $X \in \mathcal{F}_0(T_k)$.
(2) $X \cong P_k$.

Proof. Since T_k and X are direct summands of the cluster tilting module, we have $\operatorname{Ext}^i_\Lambda(T_k, X) = 0$ for any $0 < i < n$. Clearly we have $\operatorname{Ext}^i_\Lambda(T_k, X) = 0$ for any $i > n$ since $\operatorname{gl.dim}\Lambda \leq n$. On the other hand, by Proposition 2.5 (3), we have

$$\operatorname{Ext}^n_\Lambda(T_k, X) \cong \operatorname{Ext}^n_\Lambda(\tau_n^- P_k, X) \cong D\overline{\operatorname{Hom}}_\Lambda(X, P_k).$$

Assume that $X \in \mathcal{F}_0(T_k)$. Then since we have $\overline{\operatorname{Hom}}_\Lambda(X, P_k) = 0$, we have $X \not\cong P_k$. Conversely if $X \not\cong P_k$, we obtain $\overline{\operatorname{Hom}}_\Lambda(X, P_k) = 0$ since P_k is simple projective. Hence we have $X \in \mathcal{F}_0(T_k)$. □

Then by combining with Lemma 2.9 and 2.10, we have the next corollary. This is essentially shown in [IO1, Theorem 4.2].

Corollary 2.11. *Let* $\Lambda = KQ/(R)$ *be an* n*-representation-finite algebra,* $k \in Q_0$ *be a sink,* T_k *be the corresponding* n*-APR tilting* Λ*-module and put* $\Gamma := \operatorname{End}_\Lambda(T_k)$*. Denote by* M *and* M' *the basic* n*-cluster tilting modules of* $\operatorname{mod}\Lambda$ *and* $\operatorname{mod}\Gamma$*, respectively. Then, the functors* $\operatorname{Hom}_\Lambda(T_k, -)$ *and* $- \otimes_\Gamma T_k$ *induce quasi-inverse equivalences*

$$\operatorname{add}(M/P_k) \cong \operatorname{add}(M'/I_k),$$

where P_k *and* I_k *denote the simple projective* Λ*-module and the simple injective* Γ*-module associated with* k*, respectively.*

Furthermore, we recall the following definition.

Definition 2.12. Let Q be a finite, connected, acyclic quiver with $|Q_0| = l$. We call an ordering $i_1 \dots i_l$ of the vertices of Q *admissible* if for each j, the vertex i_j is a sink for the quiver obtained by removing vertices $i_1, \dots, i_{j-1}$ from Q.

Note that there exists an admissible ordering if Q is acyclic. Throughout this paper, we assume that Q has an admissible numbering $(1 \dots l)$ for simplicity.

Then, for an n-representation-finite algebra $\Lambda = KQ/(R)$, we have n-APR tilted algebras defined inductively by $\Lambda^1 := \Lambda, \Lambda^2 := \operatorname{End}_{\Lambda^1}(T_1), \dots, \Lambda^l := \operatorname{End}_{\Lambda^{l-1}}(T_{l-1})$ from Proposition 2.8, where T_i is the n-APR tilting Λ^i-module associated with i. Note that the quiver of Λ^i is also acyclic and has an admissible numbering. Explicit descriptions of n-APR tilted algebras are given in section 6.

Example 2.13. (1) Let Λ be the n-representation-finite algebra given by the following quiver with radical square zero relations.



Then we have n-APR tilted algebras as follows.



All relations are radical square zero.

(2) Let Λ be the 2-representation-finite algebra given by the following quiver with commutative relations and zero relations for each small half square.



Then we have 2-APR tilted algebras as follows.



All relations are commutative relations and zero relations for each small half square

More examples are given by [IO1].

3. Cluster-indecomposable modules and their properties

3.1. **Cluster-roots.** In this subsection we will show that the dimension vectors of cluster-indecomposable modules give roots of the Euler form. Throughout this subsection, let $\Lambda = KQ/(R)$ be an n-representation-finite algebra.

First we give the following lemma, which is well-known for $n = 1$ (see for example [ARS, VIII. Proposition 2.2]).

Lemma 3.1. *Let Φ be the Coxeter transformation and X be a cluster-indecomposable Λ-module.*

(1) *If X is non-projective, then we have $\underline{\dim}(\tau_n X) = \Phi(\underline{\dim} X)$.*

(2) *If X is non-injective, then we have* $\underline{\dim}(\tau_n^- X) = \Phi^-(\underline{\dim} X)$.

Proof. We only prove (1); the proof of (2) is similar. Since Φ gives the action of ν_n on the Grothendieck group $K_0(\mathcal{D}^b(\operatorname{mod}\Lambda))$, we have $\underline{\dim}(\nu_n X) = \Phi(\underline{\dim} X)$. On the other hand, we have $\nu_n X \cong \tau_n X$ by Proposition 2.5 (6). Thus we obtain $\underline{\dim}(\tau_n X) = \underline{\dim}(\nu_n X) = \Phi(\underline{\dim} X)$. ☐

By using Lemma 3.1, we have the following result.

Theorem 3.2. *Let $\Lambda = KQ/(R)$ be an n-representation-finite algebra and q_Λ be the Euler form of Λ. Then for any cluster-indecomposable Λ-module X, the dimension vector $\underline{\dim} X$ is a positive root of q_Λ.*

Proof. By Proposition 2.5 (1), there exist non-negative integers j and $i \in Q_0$ such that

$$X \cong \tau_n^j I_i.$$

Then, by Lemma 3.1, we have

$$\underline{\dim} X = \Phi^j(\underline{\dim} I_i).$$

Then, since Q is acyclic and by Lemma 2.2, we have

$$\begin{aligned} q_\Lambda(\underline{\dim} X) &= q_\Lambda(\Phi^j(\underline{\dim} I_i)) \\ &= q_\Lambda(\underline{\dim} I_i) \\ &= \sum_{j \geq 0} (-1)^j \dim \operatorname{Ext}^j_\Lambda(I_i, I_i) \\ &= \dim \operatorname{Hom}_\Lambda(I_i, I_i) \\ &= 1. \end{aligned}$$

Thus $\underline{\dim} X$ is a root of q_Λ. ☐

By this theorem, it is reasonable to use the following terminology.

Definition 3.3. Let Λ be an n-representation-finite algebra. We call a root $x \in \mathbb{Z}^l$ *cluster-root* if there is a cluster-indecomposable module X such that $\underline{\dim} X = x$.

By the above result and Proposition 2.5 (1), we can easily calculate cluster-roots by applying the Coxeter transformation to the dimension vectors of indecomposable injective (or projective) modules.

Example 3.4. Let Λ be the following algebra with commutative relations and zero relations for each small half square

$$\begin{tikzcd}
 & & 3 \arrow[dr] & & \\
 & 5 \arrow[ur] \arrow[dr] \arrow[rr, dash, dashed] & & 2 \arrow[dr] & \\
6 \arrow[ur] \arrow[rr, dash, dashed] & & 4 \arrow[ur] \arrow[rr, dash, dashed] & & 1.
\end{tikzcd}$$

Then the Coxeter transformation Φ is the matrix

$$\Phi = C_\Lambda^t C_\Lambda^{-1} = \begin{pmatrix} 1 & -1 & 0 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}.$$

Then the cluster-roots are given as follows

$$\underline{\dim} I_1 = \begin{pmatrix} 1\\1\\1\\0\\0\\0 \end{pmatrix}, \underline{\dim} I_2 = \begin{pmatrix} 0\\1\\1\\1\\1\\0 \end{pmatrix}, \underline{\dim} I_3 = \begin{pmatrix} 0\\0\\1\\0\\1\\1 \end{pmatrix}, \underline{\dim} I_4 = \begin{pmatrix} 0\\0\\0\\1\\1\\0 \end{pmatrix}, \underline{\dim} I_5 = \begin{pmatrix} 0\\0\\0\\0\\1\\1 \end{pmatrix}, \underline{\dim} I_6 = \begin{pmatrix} 0\\0\\0\\0\\0\\1 \end{pmatrix},$$

$$\underline{\dim} P_2 = \Phi(\underline{\dim} I_4) = \begin{pmatrix} 1\\1\\0\\0\\0\\0 \end{pmatrix}, \underline{\dim} P_4 = \Phi(\underline{\dim} I_5) = \begin{pmatrix} 0\\1\\0\\1\\0\\0 \end{pmatrix},$$

$$\underline{\dim} S_4 = \Phi(\underline{\dim} I_6) = \begin{pmatrix} 0\\0\\0\\1\\0\\0 \end{pmatrix}, \underline{\dim} P_1 = \Phi^2(\underline{\dim} I_6) = \begin{pmatrix} 1\\0\\0\\0\\0\\0 \end{pmatrix}.$$

Furthermore, as we will show in section 3, cluster-indecomposable modules are uniquely determined by their dimension vectors. Thus we can find the number of cluster-indecomposable modules and their the dimension vectors by simple calculation.

For later use, we will show that the Coxeter transformation has finite order.

Proposition 3.5. *Let Φ be the Coxeter transformation of Λ. Then there exists an integer d such that $\Phi^d = \mathbf{1}$.*

Proof. Since the global dimension of Λ is finite, the dimension vectors of the complete set $\{\underline{\dim} P_1, \ldots, \underline{\dim} P_l\}$ of the indecomposable projective modules generate the canonical basis of the free group $\mathbb{Z}^l$. Therefore it is enough to show that there exists an integer d such that $\Phi^d(\underline{\dim} P_i) = \underline{\dim} P_i$ for any $i \in Q_0$.

By [HI1, Theorem 1.1], Λ is a twisted fractionally Calabi-Yau algebra. Therefore there exist integers s, t such that $\nu_n^s \Lambda \cong \Lambda[t]$ in $\mathcal{D}^b(\mathrm{mod}\,\Lambda)$ (see [HI1, Proposition 4.2]). Thus there exists an integer d such that $\Phi^d(\underline{\dim} P_i) = \underline{\dim} P_i$ for any $i \in Q_0$. □

3.2. **Reflection transformations.** In [BGP], the authors showed that reflection functors were compatible with the action of the Weyl group in the Grothendieck group, and by using this property, they gave a conceptual proof of Gabriel's theorem. Following their results, we modify the classical reflection transformations, and show the commutativity with our reflection functors in a nice subcategory. This property plays an essential role for establishing relationships between cluster-indecomposable modules and cluster-roots.

Definition 3.6. Let $\Lambda = KQ/(R)$ be a finite dimensional algebra with finite global dimension n. Then for the Euler form $\langle -, - \rangle$ of Λ, the *symmetrized Euler form* $(-,-)$ is defined by

$$(x, y) := \frac{1}{2}[\langle x, y\rangle + \langle y, x\rangle].$$

For each $i \in Q_0$, we define $\mathbb{Z}$-linear map given by

$$s_i : \mathbb{Z}^l \to \mathbb{Z}^l,\ s_i(x) := x - 2(x, \mathbf{e}_i)\mathbf{e}_i,$$

$$\delta_i : \mathbb{Z}^l \to \mathbb{Z}^l,\ \delta_i(x) := (x_1, \ldots, x_{i-1}, (-1)^{n-1}x_i, x_{i+1}, \ldots, x_l).$$

We define $t_i := \delta_i s_i : \mathbb{Z}^l \to \mathbb{Z}^l$ and call this a *reflection transformation.*

Reflection functors are defined in terms of linear representations of quivers [BGP]. They are now formulated in terms of APR tilting modules from a functorial viewpoint [APR]. Now we define reflection functors via n-APR tilting modules as follows.

Let $\Lambda = KQ/(R)$ be an n-representation-finite algebra, $k \in Q_0$ be a sink and T_k be corresponding the n-APR tilting module and put $\Gamma := \mathrm{End}_\Lambda(T_k)$. Then we define *reflection functors*

$$\begin{aligned}\mathbf{T}_k^+ &:= \mathrm{Hom}_\Lambda(T_k, -) : \operatorname{mod}\Lambda \to \operatorname{mod}\Gamma,\\ \mathbf{T}_k^- &:= - \otimes_\Gamma T_k : \operatorname{mod}\Gamma \to \operatorname{mod}\Lambda.\end{aligned}$$

Note that we have $\mathbf{T}_k^+ X = 0$ if and only if X is isomorphic to the simple projective Λ-module P_k. Moreover if X is a cluster-indecomposable Λ-module and $X \ncong P_k$, we have $\mathbf{T}_k^- \mathbf{T}_k^+(X) \cong X$ by Corollary 2.11.

Then we give a relationship between the reflection functors and reflection transformations as follows. Here we use the notations $\mathcal{F}_0, \mathcal{X}_0$ as defined in section 2.3.

Proposition 3.7. *Let $\Lambda = KQ/(R)$ be an n-representation-finite algebra, $k \in Q_0$ be a sink and T_k be the corresponding n-APR tilting module. Put $\Gamma := \mathrm{End}_\Lambda(T_k)$.*

1. *Let X be an indecomposable Λ-module. If $X \in \mathcal{F}_0(T_k)$, then $\mathbf{T}_k^+ X$ is an indecomposable Γ-module and we have $\underline{\dim}(\mathbf{T}_k^+ X) = t_k(\underline{\dim} X)$.*
2. *Let Y be an indecomposable Γ-module. If $Y \in \mathcal{X}_0(T_k)$, then $\mathbf{T}_k^- Y$ is an indecomposable Λ-module and we have $\underline{\dim}(\mathbf{T}_k^- Y) = t_k^{-1}(\underline{\dim} Y)$.*

Proof. We only prove (1); the proof of (2) is similar. Since the functor $\mathbf{T}_k^+$ gives an equivalence between $\mathcal{F}_0(T_k)$ and $\mathcal{X}_0(T_k)$ by Lemma 2.9, $\mathbf{T}_k^+ X$ is an indecomposable Γ-module. We denote by $(\underline{\dim} X)_i$ the i-th coordinate of the dimension vector $\underline{\dim} X$ for $i \in Q_0$. Assume that $i \neq k$. Then by Lemma 2.9, we have

$$\begin{aligned}(\underline{\dim}(\mathbf{T}_k^+ X))_i &= \dim \mathrm{Hom}_\Gamma(e_i\Gamma, \mathbf{T}_k^+ X)\\ &= \dim \mathrm{Hom}_\Gamma(\mathrm{Hom}_\Lambda(T_k, e_i\Lambda), \mathrm{Hom}_\Lambda(T_k, X))\\ &= \dim \mathrm{Hom}_\Lambda(e_i\Lambda, X)\\ &= (\dim X)_i\\ &= (t_k(\dim X))_i.\end{aligned}$$

If $i = k$, we have

$$\begin{aligned}(\mathbf{T}_k^+ X)e_k &\cong \mathrm{Hom}_\Gamma(e_k\Gamma, \mathbf{T}_k^+ X)\\ &\cong \mathrm{Hom}_\Gamma(\mathrm{Hom}_\Lambda(T_k, \tau_n^- P_k), \mathrm{Hom}_\Lambda(T_k, X))\\ &\cong \mathrm{Hom}_\Lambda(\tau_n^- P_k, X).\end{aligned}$$

On the other hand, we have

$$\begin{aligned}(s_k(\underline{\dim} X))_k &= \dim X e_k - [\langle \underline{\dim} X, \mathbf{e}_k\rangle + \langle \mathbf{e}_k, \underline{\dim} X\rangle]\\ &= -\langle \underline{\dim} X, \mathbf{e}_k\rangle.\end{aligned}$$

Then, by Lemma 2.2 and Lemma 3.1, we obtain

$$\begin{aligned}\langle \underline{\dim} X, \mathbf{e}_k\rangle &= (-1)^n \langle \Phi^{-1}(\mathbf{e}_k), \underline{\dim} X\rangle\\ &= (-1)^n \langle \underline{\dim}(\tau_n^- P_k), \underline{\dim} X\rangle\\ &= (-1)^n \sum_{j\geq 0} (-1)^j \dim \mathrm{Ext}_\Lambda^j(\tau_n^- P_k, X).\end{aligned}$$

Since $X \in \mathcal{F}_0(T_k)$, we have $\sum_{j\geq 0}(-1)^j \operatorname{Ext}^j_\Lambda(\tau_n^- P_k, X) = \operatorname{Hom}_\Lambda(\tau_n^- P_k, X)$. Hence we obtain

$$\begin{aligned}
(t_k(\underline{\dim} X))_k &= (-1)^{n-1}(s_k(\underline{\dim} X))_k \\
&= (-1)^{n-1}(-\langle \underline{\dim} X, \mathbf{e}_k\rangle) \\
&= (-1)^{n-1}\{(-1)(-1)^n \dim \operatorname{Hom}_\Lambda(\tau_n^- P_k, X)\} \\
&= \dim \operatorname{Hom}_\Lambda(\tau_n^- P_k, X).
\end{aligned}$$

Thus we have shown that $\underline{\dim}(\mathbf{T}_k^+ X) = t_k(\underline{\dim} X)$. □

As an immediate consequence, we have the following.

Corollary 3.8. *Let $\Lambda = KQ/(R)$ be an n-representation-finite algebra, $k \in Q_0$ be a sink and T_k be the n-APR tilting module. Put $\Gamma := \operatorname{End}_\Lambda(T_k)$.*

1. *Let X be a cluster-indecomposable Λ-module. If $X \ncong P_k$, then $\mathbf{T}_k^+ X$ is an indecomposable Γ-module and $\underline{\dim}(\mathbf{T}_k^+ X) = t_1(\underline{\dim} X)$.*
2. *Let Y be a cluster-indecomposable Γ-module. If $Y \ncong I_k$, then $\mathbf{T}_k^- Y$ is an indecomposable Λ-module and $\underline{\dim}(\mathbf{T}_k^- Y) = t_1^-(\underline{\dim} Y)$.*

Proof. We only prove (1); the proof (2) is similar. By Proposition 3.7, it is enough to show $X \in \mathcal{F}_0(T_k)$. We have shown this in Lemma 2.10. □

Next we deal with $n = 2$ for later use. We note that, if $\operatorname{gl.dim} \Lambda \leq 2$, then q_Λ is given by the *Tits form* [B2] and it can be characterized in terms of a quiver with relations.

Lemma 3.9. *Let $\Lambda = KQ/(R)$ be a 2-representation-finite algebra with a minimal set R of relations, $k \in Q_0$ be a sink and T_k be the 2-APR tilting module. Put $\Gamma := \operatorname{End}_\Lambda(T_k)$.*

1. *We have $q_\Gamma(x) = q_\Lambda(\delta_k x)$.*
2. *If x is a root of q_Λ , then $t_k x$ is a root of q_Γ.*

Proof. Since $\operatorname{gl.dim} \Lambda \leq 2$ and $\operatorname{gl.dim} \Gamma \leq 2$, we can describe q_Λ and q_Γ by the Tits forms. Then because k is sink, we have

$$q_\Lambda(x) = \sum_{i \in Q_0} x_i^2 - \sum_{\substack{a \in Q_1 \\ e(a)=k}} x_{s(a)} x_k + \sum_{\substack{r \in R \\ e(r)=k}} x_{s(r)} x_k + \alpha,$$

where α has no term containing x_k. Let (Q', R') be the quiver with a minimal set of relations satisfying $KQ'/(R') \cong \Gamma$. Then by [IO1, Theorem 3.11] (cf. Proposition 6.3), we have

$$\begin{aligned}
q_\Gamma(x) &= \sum_{i \in Q_0} x_i^2 + \sum_{\substack{r \in R' \\ s(r)=k}} x_{e(r)} x_k - \sum_{\substack{a \in Q_1' \\ s(a)=k}} x_{e(a)} x_k + \alpha \\
&= \sum_{i \in Q_0} x_i^2 - \sum_{\substack{a \in Q_1 \\ e(a)=k}} x_{s(a)}(-x_k) + \sum_{\substack{r \in R \\ e(r)=k}} x_{s(r)}(-x_k) + \alpha.
\end{aligned}$$

Thus the first statement follows. Moreover it is well-known that $q_\Lambda(s_i x) = q_\Lambda(x)$ for any $i \in Q_0$. Then, by the first assertion, we obtain $q_\Gamma(t_k x) = q_\Lambda(\delta_k t_k x) = q_\Lambda(s_k x) = q_\Lambda(x) = 1$.

□

Example 3.10. Let Λ be the algebra of Example 3.4 and put $\Gamma := \mathrm{End}_\Lambda(T_1)$ (the quiver and relations are given in Example 2.13 (2)). We have the Euler form of Λ

$$q_\Lambda(x) = \sum_{i \in Q_0} x_i^2 - x_1x_2 - x_2x_3 - x_2x_4 - x_3x_5 - x_4x_5 - x_5x_6 + x_1x_4 + x_2x_5 + x_4x_6.$$

Then since $q_\Gamma(x) = q_\Lambda(\delta_1 x)$, we have the Euler form of Γ

$$q_\Gamma(x) = \sum_{i \in Q_0} x_i^2 + x_1x_2 - x_2x_3 - x_2x_4 - x_3x_5 - x_4x_5 - x_5x_6 - x_1x_4 + x_2x_5 + x_4x_6.$$

For example, $\mathbf{e}_4$ is one of the roots of $q_\Lambda(x)$. Then for the reflection transformation $t_1 = \begin{pmatrix} 1 & -1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$, $t_1\mathbf{e}_4$ is also a root of $q_\Gamma(x)$.

4. Cluster-roots determine cluster-indecomposable modules

In this section we investigate relationships between cluster-indecomposable modules and cluster-roots in more detail. In particular, we show that cluster-indecomposable modules can be obtained from simple modules by applying a sequence of the reflection functors. As a corollary, we give a complete description of cluster roots in terms of reflection transformations. We prove that cluster-indecomposable modules are uniquely determined by their dimension vectors.

Throughout this section, let $\Lambda = KQ/(R)$ be an n-representation-finite algebra with an admissible numbering $(1 \ldots l)$. As we have seen, we have n-APR tilted algebras $\Lambda^1 := \Lambda, \Lambda^2 := \mathrm{End}_{\Lambda^1}(T_1), \ldots, \Lambda^{l+1} := \mathrm{End}_{\Lambda^l}(T_l)$. We denote by t_i the reflection transformation over Λ^i and denote by $P_j^{(i)}$ the Λ^i-module P_j for $i \in Q_0$. We write

$$\begin{aligned} \mathbf{T}_i^+ &:= \mathrm{Hom}_{\Lambda^i}(T_i, -) : \mathrm{mod}\,\Lambda^i \to \mathrm{mod}\,\Lambda^{i+1}, \\ \mathbf{T}_i^- &:= - \otimes_{\Lambda^{i+1}} T_i : \mathrm{mod}\,\Lambda^{i+1} \to \mathrm{mod}\,\Lambda^i, \end{aligned}$$

and we write

$$\begin{aligned} \mathbb{R}\mathbf{T}_i^+ &:= \mathbb{R}\mathrm{Hom}_{\Lambda^i}(T_i, -) : \mathcal{D}^b(\mathrm{mod}\,\Lambda^i) \to \mathcal{D}^b(\mathrm{mod}\,\Lambda^{i+1}), \\ \mathbb{L}\mathbf{T}_i^- &:= - \overset{\mathbb{L}}{\otimes}_{\Lambda^{i+1}} T_i : \mathcal{D}^b(\mathrm{mod}\,\Lambda^{i+1}) \to \mathcal{D}^b(\mathrm{mod}\,\Lambda^i). \end{aligned}$$

First we give the following easy observation.

Lemma 4.1. *We have isomorphisms*

$$\mathbb{R}\mathbf{T}_i^+(\nu_n^- P_j^{(i)}) \cong \begin{cases} \nu_n^- P_j^{(i+1)} & (j \neq i) \\ P_j^{(i+1)} & (j = i), \end{cases}$$

$$\mathbb{L}\mathbf{T}_i^-(\nu_n^- P_j^{(i+1)}) \cong \begin{cases} \nu_n^- P_j^{(i)} & (j \neq i) \\ \nu_n^-(\tau_n^- P_j^{(i)}) & (j = i). \end{cases}$$

Proof. We only prove the first statement; the proof of the second one is similar. By the definition, we have $P_j^{(i+1)} = \mathrm{Hom}_{\Lambda^i}(T_i, P_j^{(i)})$ for $j \neq i$ and $P_i^{(i+1)} = \mathrm{Hom}_{\Lambda^i}(T_i, \tau_n^- P_i^{(i)})$.

Assume that $j \neq i$. Then we have

$$\begin{aligned} \mathbb{R}\mathbf{T}_i^+(\nu_n^- P_j^{(i)}) &\cong \nu_n^-(\mathbb{R}\mathbf{T}_i^+(P_j^{(i)})) \\ &\cong \nu_n^-(\mathbf{T}_i^+(P_j^{(i)})) \quad (\mathrm{Ext}^m_{\Lambda^i}(T_i, P_j^{(i)}) = 0, m > 0) \\ &\cong \nu_n^-(P_j^{(i+1)}). \end{aligned}$$

Assume that $j = i$. Then since $P_i^{(i)}$ is not an injective Λ^i-module, we have $\nu_n^- P_j^{(i)} \cong \tau_n^- P_j^{(i)}$ by Proposition 2.5 (6). Thus we obtain

$$\begin{aligned} \mathbb{R}\mathbf{T}_i^+(\nu_n^- P_j^{(i)}) &\cong \mathbb{R}\mathbf{T}_i^+(\tau_n^- P_j^{(i)}) \\ &\cong \mathbf{T}_i^+(\tau_n^- P_j^{(i)}) \quad (\mathrm{Ext}^m_{\Lambda^i}(T_i, \tau_n^- P_j^{(i)}) = 0, m > 0) \\ &\cong P_j^{(i+1)}. \end{aligned}$$

□

Then we give the following fundamental property of n-representation-finite algebras. This proposition explains the reason that the algebras return to the original algebras by n-APR tilts in example 2.13.

Proposition 4.2. *We have an isomorphism* $\Lambda^{l+1} \cong \Lambda$.

Proof. By applying Lemma 4.1 repeatedly, we obtain $\Lambda^{l+1} \cong \mathbb{R}\mathbf{T}_l^+ \ldots \mathbb{R}\mathbf{T}_1^+(\nu_n^- \Lambda)$. Then since $\mathbb{R}\mathbf{T}_i^+$ and ν_n^- are equivalences, we obtain

$$\begin{aligned} \Lambda^{l+1} &\cong \mathrm{End}_{\Lambda^{l+1}}(\Lambda^{l+1}) \\ &\cong \mathrm{End}_{\mathcal{D}(\Lambda)}(\mathbb{R}\mathbf{T}_l^+ \ldots \mathbb{R}\mathbf{T}_1^+(\nu_n^- \Lambda)) \\ &\cong \mathrm{End}_{\mathcal{D}(\Lambda)}(\nu_n^- \Lambda) \\ &\cong \mathrm{End}_{\mathcal{D}(\Lambda)}(\Lambda) \\ &\cong \Lambda. \end{aligned}$$

□

The next lemma is analogous to [APR, Proposition 2.6], which gives a relationship between the indecomposable injective modules over Λ^i and Λ^{i+1}.

Lemma 4.3. *Fix a vertex* $i \in Q_0$. *Let* $\{I_1, \ldots, I_l\}$ *be a complete set of the indecomposable injective* Λ^i*-modules. Then* $\{\mathbf{T}_i^+(I_1), \ldots, \mathbf{T}_i^+(I_{i-1}), \mathrm{Ext}^n_\Lambda(T_i, P_i), \mathbf{T}_i^+(I_{i+1}), \ldots, \mathbf{T}_i^+(I_l)\}$ *is a complete set of the indecomposable injective* Λ^{i+1}*-modules.*

Proof. Let $j \in Q_0$ be a vertex with $j \neq i$. Then we obtain Λ^{i+1}-module isomorphisms

$$\begin{aligned} \mathbf{T}_i^+(I_j) &\cong \mathrm{Hom}_{\Lambda^i}(T_i, D(\Lambda^i e_j)) \\ &\cong D(T_i \otimes_{\Lambda^i} \Lambda^i e_j) \\ &\cong D((T_i)e_j). \end{aligned}$$

Moreover, we get

$$\begin{aligned} (T_i)e_j &\cong \mathrm{Hom}_{\Lambda^i}(e_j\Lambda^i, T_i) \\ &\cong \mathrm{Hom}_{\Lambda^{i+1}}(\mathrm{Hom}_{\Lambda^i}(T_i, e_j\Lambda^i), \mathrm{Hom}_{\Lambda^i}(T_i, T_i)) \\ &\cong \mathrm{Hom}_{\Lambda^{i+1}}(e_j\Lambda^{i+1}, \Lambda^{i+1}) \\ &\cong \Lambda^{i+1}e_j. \end{aligned}$$

Thus we obtain $\mathbf{T}_i^+(I_j) \cong D(\Lambda^{i+1}e_j)$ as Λ^{i+1}-modules. Then, together with Proposition 2.8, the statement follows. □

Then we give the descriptions of all projective modules and injective modules in terms of reflection functors as follows.

Proposition 4.4. (1) *Let P_i be a projective Λ-module. Then P_i is isomorphic to $\mathbf{T}_1^- \dots \mathbf{T}_{i-1}^-(S_i)$, where S_i is the simple projective Λ^i-module.*

(2) *Let I_j be a injective Λ-module. Then I_j is isomorphic to $\mathbf{T}_l^+ \dots \mathbf{T}_{j+1}^+(S_j)$, where S_j is the simple injective Λ^{j+1}-module.*

Proof. (1) Let $h \in Q_0$ be a vertex such that $h < i$. Then since we have $P_i^{(h)} = \mathrm{Hom}_{\Lambda^{h-1}}(T_{h-1}, P_i^{(h-1)}) = \mathbf{T}_{h-1}^+(P_i^{(h-1)})$, we have $\mathbf{T}_{h-1}^- P_i^{(h)} \cong \mathbf{T}_{h-1}^-(\mathbf{T}_{h-1}^+(P_i^{(h-1)})) \cong P_i^{(h-1)}$ by Corollary 2.11. Applying this process repeatedly for any $h < i$, we have the assertion.

(2) By Proposition 4.2, we have $\Lambda^{l+1} \cong \Lambda$. Then the proof is similar to that of (1) by Lemma 4.3.

□

As a corollary, we will show that the Coxeter transformation can be described as a composition of reflection transformations.

Corollary 4.5. *Let Φ be the Coxeter transformation. Then we have $\Phi = t_l \dots t_1$.*

Proof. It is enough to show that $\Phi(\underline{\dim} P_i) = t_l \dots t_1(\underline{\dim} P_i)$ for any $i \in Q_0$. By Lemma 2.2, we have

$$\Phi(\underline{\dim} P_i) = (-1)^n \underline{\dim} I_i.$$

On the other hand, by Corollary 3.8 and Proposition 4.4, we have

$$\begin{aligned} t_l \dots t_1(\underline{\dim} P_i) &= t_l \dots t_1(t_1^{-1} \dots t_{i-1}^{-1}\mathbf{e}_i) \\ &= t_l \dots t_i \mathbf{e}_i \\ &= (-1)^n t_l \dots t_{i+1}\mathbf{e}_i \\ &= (-1)^n \underline{\dim} I_i. \end{aligned}$$

□

Next we give a relationship between reflection functors and n-AR translations.

Lemma 4.6. *We have an isomorphism $\tau_n^- \Lambda \cong \mathbf{T}_1^- \dots \mathbf{T}_{l-1}^-(T_l)$, where T_l is the n-APR tilting Λ^l-module.*

Proof. We denote by $U_i := (\bigoplus_{j=1}^{i-1} P_j^{(i)}) \oplus \tau_n^-(\bigoplus_{j=i}^{l} P_j^{(i)})$. Note that we have $U_1 = \tau_n^- \Lambda$ and $U_l = T_l$. Then it is enough to show that

$$U_{i-1} \cong \mathbf{T}_{i-1}^- U_i,$$

for any $i \in Q_0$ with $i > 1$. First, for a vertex $j \in Q_0$ such that $j < i-1$, we have $P_j^{(i-1)} \cong \mathbf{T}_{i-1}^- P_j^{(i)}$ by Corollary 2.11. Similarly, for $j = i-1$, we have $\tau_n^- P_j^{(i-1)} \cong \mathbf{T}_{i-1}^- P_j^{(i)}$.

Next, let $j \in Q_0$ be a vertex such that $j \geq i$.

(i) Assume that $P_j^{(i)}$ is not injective. Then since we obtain $\tau_n^- P_j^{(i)} \cong \nu_n^- P_j^{(i)}$ by Proposition 2.5 (6), we have

$$\begin{aligned}
\mathbf{T}_{i-1}^-(\tau_n^- P_j^{(i)}) &\cong H^0(\mathbb{L}\mathbf{T}_{i-1}^-(\nu_n^- P_j^{(i)})) \\
&\cong H^0(\nu_n^- P_j^{(i-1)}) \qquad \text{(Lemma 4.1)} \\
&\cong \tau_n^- P_j^{(i-1)}.
\end{aligned}$$

(ii) Assume that $P_j^{(i)}$ is injective. Then by Lemma 4.3, $P_j^{(i-1)}$ is also injective. Therefore we have $\tau_n^- P_j^{(i)} = 0$ and $\tau_n^- P_j^{(i-1)} = 0$.

Consequently, we get $\tau_n^- P_j^{(i-1)} \cong \mathbf{T}_{i-1}^-(\tau_n^- P_j^{(i)})$ for $j \geq i$. Thus we obtain $U_{i-1} \cong \mathbf{T}_{i-1}^- U_i$ for $i \in Q_0$ and the statement follows from the inductive step. ☐

Now we define the *Coxeter functors* by

$$\begin{aligned}
\mathbf{C}^+ &:= \mathbf{T}_l^+ \dots \mathbf{T}_1^+ : \operatorname{mod}\Lambda \to \operatorname{mod}\Lambda, \\
\mathbf{C}^- &:= \mathbf{T}_1^- \dots \mathbf{T}_l^- : \operatorname{mod}\Lambda \to \operatorname{mod}\Lambda.
\end{aligned}$$

Then we explain the interplay between n-AR translations and the Coxeter functors. We remark that $\mathbf{C}^+, \mathbf{C}^-$ do not depend on the choice of an admissible numbering by the following result, which is an analogous result of [APR].

Proposition 4.7. (1) *There exists an isomorphism $\mathbf{C}^+X \cong \tau_n X$ for a Λ-module X.*
(2) *There exists an isomorphism $\mathbf{C}^-X \cong \tau_n^- X$ for a Λ-module X.*

Proof. (1) We have isomorphisms

$$\begin{aligned}
\mathbf{T}_l^+ \dots \mathbf{T}_1^+(X) &\cong \operatorname{Hom}_{\Lambda^l}(T_l, \operatorname{Hom}_{\Lambda^{l-1}}(T_{l-1}, \cdots (\operatorname{Hom}_\Lambda(T_1, X) \cdots) \\
&\cong \operatorname{Hom}_\Lambda(T_l \otimes_{\Lambda^l} \dots \otimes_{\Lambda^2} T_1, X) \\
&\cong \operatorname{Hom}_\Lambda(\mathbf{T}_1^- \dots \mathbf{T}_{l-1}^-(T_l), X).
\end{aligned}$$

Then, by Lemma 4.6, we have $\mathbf{C}^+X \cong \operatorname{Hom}_\Lambda(\tau_n^-\Lambda, X)$.

Moreover we have $\operatorname{Hom}_\Lambda(\tau_n^-\Lambda, \Lambda) = 0$ by Proposition 2.5 (4). Thus we get $\operatorname{Hom}_\Lambda(\tau_n^-\Lambda, X) \cong \underline{\operatorname{Hom}}_\Lambda(\tau_n^-\Lambda, X)$. Then, using Proposition 2.5 (3), we obtain

$$\begin{aligned}
\operatorname{Hom}_\Lambda(\tau_n^-\Lambda, X) &\cong \underline{\operatorname{Hom}}_\Lambda(\tau_n^-\Lambda, X) \\
&\cong D\operatorname{Ext}_\Lambda^n(X, \tau_n(\tau_n^-\Lambda)) \\
&\cong D\operatorname{Ext}_\Lambda^n(X, \Lambda/I) \quad (I : \text{injective modules of } \Lambda) \\
&\cong D\operatorname{Ext}_\Lambda^n(X, \Lambda) \\
&= \tau_n X.
\end{aligned}$$

(2) We have isomorphisms

$$\begin{aligned}
D(\mathbf{T}_1^- \dots \mathbf{T}_l^-(X)) &\cong \operatorname{Hom}_K(\mathbf{T}_1^- \dots \mathbf{T}_l^-(X), K) \\
&\cong \operatorname{Hom}_K(X \otimes_{\Lambda^{l+1}} T_l \otimes_{\Lambda^l} \dots \otimes_{\Lambda^2} T_1, K) \\
&\cong \operatorname{Hom}_\Lambda(X, \operatorname{Hom}_K(T_l \otimes_{\Lambda^l} \dots \otimes_{\Lambda^2} T_1, K)) \\
&\cong \operatorname{Hom}_\Lambda(X, \operatorname{Hom}_K(\mathbf{T}_1^- \dots \mathbf{T}_{l-1}^-(T_l), K)).
\end{aligned}$$

By Lemma 4.6, we have $D(\mathbf{C}^-X) \cong \operatorname{Hom}_\Lambda(X, D(\tau_n^-\Lambda)) \cong \operatorname{Hom}_\Lambda(X, \tau_n(D\Lambda))$.

Then, similarly we obtain

$$\begin{aligned} D\operatorname{Hom}_\Lambda(X,\tau_n(D\Lambda)) &\cong D\overline{\operatorname{Hom}}_\Lambda(X,\tau_n(D\Lambda)) \\ &\cong \operatorname{Ext}^n_\Lambda(\tau_n^-(\tau_n(D\Lambda)),X) \\ &\cong \operatorname{Ext}^n_\Lambda((D\Lambda)/P,X) \quad (P : \text{projective modules of } D\Lambda) \\ &\cong \operatorname{Ext}^n_\Lambda(D\Lambda,X) \\ &= \tau_n^- X. \end{aligned}$$

□

Thus if X is projective (injective, respectively), then we obtain $\mathbf{C}^+X \cong \tau_n X = 0$ ($\mathbf{C}^-X \cong \tau_n^- X = 0$). The reason also follows from Proposition 4.4.

Consequently, we show that any cluster-indecomposable module can be obtained from a simple module by applying a sequence of reflection functors. This is a generalization of [BGP, Corollary 3.1].

Corollary 4.8. *Let X be a cluster-indecomposable Λ-module.*

1. *There exists an integer $u \geq 0$ and $i \in Q_0$ such that $X \cong \mathbf{C}^{-u}\mathbf{T}_1^- \dots \mathbf{T}_{i-1}^-(S_i)$, where S_i is the simple projective Λ^i-module.*
2. *There exists an integer $v \geq 0$ and $j \in Q_0$ such that $X \cong \mathbf{C}^{v}\mathbf{T}_l^+ \dots \mathbf{T}_{j+1}^+(S_j)$, where S_j is the simple injective Λ^{j+1}-module.*

Proof. We only prove (1); the proof of (2) is similar. By Proposition 2.5 (5) and Proposition 4.7, there exists an integer $u \geq 0$ such that $\mathbf{C}^uX \neq 0$ and $\mathbf{C}^{u+1}X = 0$. Hence there exists $i \in Q_0$ such that $\mathbf{T}_{i-1}^+\mathbf{T}_{i-2}^+ \dots \mathbf{T}_1^+\mathbf{C}^u(X) \neq 0$ and $\mathbf{T}_i^+\mathbf{T}_{i-1}^+ \dots \mathbf{T}_1^+\mathbf{C}^u(X) = 0$.

Thus, we obtain $\mathbf{T}_{i-i}^+ \dots \mathbf{T}_1^+\mathbf{C}^u(X) \cong S_i$, where S_i is the simple projective Λ^i-module. By applying Corollary 2.11 repeatedly, we have $X \cong \mathbf{C}^{-u}\mathbf{T}_1^- \dots \mathbf{T}_{i-1}^-(S_i)$. □

Remark 4.9. By Proposition 4.4, $\mathbf{T}_1^- \dots \mathbf{T}_{i-1}^-(S_i)$ ($\mathbf{T}_l^+ \dots \mathbf{T}_{j+1}^+(S_j)$, respectively) is a projective module (injective module). On the other hand, the Coxeter functors $\mathbf{C}^+, \mathbf{C}^-$ give n-AR translations τ_n, τ_n^- by Proposition 4.7. Thus Corollary 4.8 rephrases Proposition 2.5 (1) in terms of reflection functors.

As a corollary of the previous results, we give the descriptions of cluster-roots in terms of reflection transformations as follows.

Theorem 4.10. *We denote by $\mathbf{p}_i = t_1^{-1} \dots t_{i-1}^{-1}\mathbf{e}_i, \mathbf{q}_j = t_l \dots t_{j+1}\mathbf{e}_j$ and $\mathbf{c} = t_l \dots t_1$.*

1. *Any cluster-root can be written as $\mathbf{c}^{-u}\mathbf{p}_i$ for an integer $u \geq 0$ and $i \in Q_0$.*
2. *Any cluster-root can be written as $\mathbf{c}^{v}\mathbf{q}_j$ for an integer $v \geq 0$ and $j \in Q_0$.*

Proof. We only prove (1); the proof (2) is similar. Let x be a cluster-root. By definition, we can take a cluster-indecomposable module X such that $\underline{\dim}X = x$. By Corollary 4.8, there exists an integer $u \geq 0$ and $i \in Q_0$ such that $X \cong \mathbf{C}^{-u}\mathbf{T}_1^- \dots \mathbf{T}_{i-1}^-(S_i)$. Then, applying Corollary 3.8 repeatedly, we obtain $\underline{\dim}X = \mathbf{c}^{-u}t_1^- \dots t_{i-1}^-(\underline{\dim}S_i) = \mathbf{c}^{-u}\mathbf{p}_i$. □

Example 4.11. Let Λ be the algebra of Example 3.4 (the quiver with relations of Λ^i is given in Example 2.13 (2)). Then we have

$$t_1^- = \begin{pmatrix} 1 & 1 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, t_2^- = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & 1 & 1 & -1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, t_3^- = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

$$t_4^- = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & -1 & 0 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, t_5^- = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, t_6^- = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & -1 & 1 \end{pmatrix}.$$

Thus we have

$$\mathbf{c}^- = t_1^- \dots t_l^- = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & -1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 & 1 \end{pmatrix}.$$

Applying $\mathbf{c}^-$ to $\mathbf{p}_i$, we have all cluster-roots as follows.

$$\mathbf{p}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 1 \\ 1 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \mathbf{p}_1,$$

$$\mathbf{p}_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 1 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \begin{pmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 1 \\ 0 \end{pmatrix} \xrightarrow{\mathbf{c}^-} \mathbf{p}_2.$$

Here we have $\mathbf{p}_3 = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$, $\mathbf{p}_4 = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}$, $\mathbf{p}_5 = \begin{pmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 1 \\ 0 \end{pmatrix}$, $\mathbf{p}_6 = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 1 \\ 1 \end{pmatrix}$.

We can check that $\mathbf{c}^-$ and $\mathbf{p}_i$ coincide with Φ^- and $\underline{\dim} P_i$, respectively. This follows from the results of Proposition 4.4 and Corollary 4.5.

Finally, we give the following theorem.

Theorem 4.12. *Let $\Lambda = KQ/(R)$ be an n-representation-finite algebra. Cluster-indecomposable modules are uniquely determined up to isomorphism by their dimension vectors.*

Proof. Assume that X and Y are cluster-indecomposable Λ-modules such that $\underline{\dim} X = \underline{\dim} Y$. By Corollary 4.8, there exists an integer $u \geq 0$ and $i \in Q_0$ such that $\mathbf{T}_{i-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^u(X) \cong S_i$, where S_i is the simple projective Λ^i-module. Note that u, i are minimal integers with this property, that is, $\mathbf{T}_{j-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X)$ is not isomorphic to the simple projective module S_j if $vl + j < ul + i$. By Corollary 3.8, we have

$$\begin{aligned} \underline{\dim}(\mathbf{T}_{i-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^u(X)) &= t_{i-1} \dots t_1 \mathbf{c}^u (\underline{\dim} X) \\ &= t_{i-1} \dots t_1 \mathbf{c}^u (\underline{\dim} Y) \\ &= \mathbf{e}_i, \end{aligned}$$

where $\mathbf{c} = t_l \dots t_1$. Thus we have $\mathbf{e}_i = t_{i-1} \dots t_1 \mathbf{c}^u (\underline{\dim} Y)$.

Then we obtain that $S_i \cong \mathbf{T}^+_{i-1} \dots \mathbf{T}^+_1 \mathbf{C}^u(Y)$. Indeed, by the minimality of u, i, $\mathbf{T}^+_{j-1} \dots \mathbf{T}^+_1 \mathbf{C}^v Y$ is cluster-indecomposable if $vl + j < ul + i$ by Corollary 2.11 and the assertion holds. Thus, we have

$$\mathbf{T}^+_{i-1} \dots \mathbf{T}^+_1 \mathbf{C}^u(X) \cong S_i \cong \mathbf{T}^+_{i-1} \dots \mathbf{T}^+_1 \mathbf{C}^u(Y).$$

Again by Corollary 2.11, we obtain $X \cong \mathbf{C}^{-u} \mathbf{T}^-_1 \dots \mathbf{T}^-_{i-1}(S_i) \cong Y$. □

5. A CRITERION FOR CLUSTER-ROOTS

In previous sections, we investigate some properties of cluster-roots. As we can observe in Example 5.4, not all positive roots are cluster-roots. Therefore it is important to characterize cluster-roots. We will show in Theorem 5.2 that cluster-roots have the Φ-sign-coherent property and, if n is even, cluster-roots are Φ-positive. Then we conjecture that the converse is true for $n = 2$. We remark that the structure of 2-representation-finite algebras is investigated in [HI2] (see also [IO2]).

First, we introduce the following notion.

Definition 5.1. Let Λ be an n-representation-finite algebra and Φ be the Coxeter transformation of Λ. We call $x \in \mathbb{Z}^l$ *Φ-sign-coherent* if $\Phi^m(x) \in \mathbb{Z}^l_{\geq 0}$ or $\Phi^m(x) \in \mathbb{Z}^l_{\leq 0}$ for any $m \in \mathbb{Z}$ and call $x \in \mathbb{Z}^l$ *Φ-positive* if $\Phi^m(x) \in \mathbb{Z}^l_{\geq 0}$ for any $m \in \mathbb{Z}$.

By Proposition 3.5, the Coxeter transformation has finite order. Therefore Φ-sign-coherent (Φ-positive) property can be checked by finitely many calculations.

Then we have the following consequence.

Theorem 5.2. *Let Λ be an n-representation-finite algebra. Then cluster-roots are Φ-sign-coherent. Moreover, if n is even, then cluster-roots are Φ-positive.*

Proof. For any non-projective cluster-indecomposable X, we obtain $\Phi(\underline{\dim} X) = \underline{\dim}(\tau_n X)$ by Lemma 3.1. Thus, we have $\Phi(\underline{\dim} X) \in \mathbb{Z}^l_{\geq 0}$. On the other hand, for an indecomposable projective module P_i, we have $\Phi(\underline{\dim} P_i) = (-1)^n \underline{\dim} I_i$ by Lemma 2.2. Thus the statement follows. □

Now we pose the following conjecture.

Conjecture 5.3. *Let Λ be a 2-representation-finite algebra. Then Φ-positive roots are cluster-roots.*

In this section, we consider this conjecture.

Example 5.4. Let Λ be the algebra of Example 3.4. Then the Euler form q_Λ was given in Example 3.10 and all positive roots of q_Λ are given as follows

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix},$$

$$\begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}.$$

We can check that Φ-positive roots are given as follows

$$\begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 1 \\ 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

Then we can verify that they coincide with cluster-roots given in Example 3.4.

Remark 5.5. It is known that, for a Dynkin quiver, all roots are either non-negative roots or non-positive roots (see for example [ASS, VII. Lemma 4.8]). Hence all roots are Φ-sign-coherent, and positive roots correspond bijectively with the indecomposable modules by Gabriel's theorem.

Remark 5.6. For the case $n > 2$, the Φ-positive property is not enough to characterize cluster-roots. For example, let Λ be the 4-representation-finite algebra given in Example 2.13 (1). Then one can check that $\underline{\dim} S_3 = \mathbf{e}_3$ is a Φ-positive root of q_Λ, while it is not a cluster-root.

Throughout this section, unless otherwise specified, let $\Lambda = KQ/(R)$ be a 2-representation-finite algebra with an admissible numbering $(1 \ldots l)$ and a minimal set R of relations. We denote by $\Lambda^{i+1} := \operatorname{End}_{\Lambda^i}(T_i)$ the 2-APR tilted algebras for $i \in Q_0$.

The aim of this section is to show the following theorem.

Theorem 5.7. *Let Λ be a 2-representation-finite algebra. If Λ^i is representation-directed (Definition 5.13) for any $i \in Q_0$, then Conjecture 5.3 is true.*

As a corollary, we give the following.

Corollary 5.8. *Let Λ be a 2-representation-finite algebra. If Λ is iterated tilted, then Conjecture 5.3 is true.*

In this case, using Bongartz' theorem (Theorem 5.14), we have the following correspondence.

$$\begin{tikzcd}
\{\text{Indecomposable modules}\} \arrow[r, leftrightarrow, "1-1"] & \{\text{Positive roots}\} \\
\{\text{Cluster-indecomposable modules}\} \arrow[u, phantom, "\bigcup"] \arrow[r, leftrightarrow, "1-1"] & \{\Phi\text{-positive roots}\} \arrow[u, phantom, "\bigcup"]
\end{tikzcd}$$

We give some properties of Φ-positive roots. We start with the following easy lemma.

Lemma 5.9. *Let Λ be a 2-representation-finite algebra, Φ be the Coxeter transformation and t_i be the reflection transformation of Λ^i for $i \in Q_0$. For $x \in \mathbb{Z}^l$, the following conditions are equivalent.*

(1) *x is Φ-positive.*
(2) *$t_i \ldots t_1 \Phi^m(x) \in \mathbb{Z}^l_{\geq 0}$ for any $i \in Q_0$ and $m \in \mathbb{Z}$.*

Proof. Clearly, (2) implies (1). Assume that $t_i \ldots t_1 \Phi^m(x) \notin \mathbb{Z}^l_{\geq 0}$ for some $i \in Q_0$ and $m \in \mathbb{Z}$. Then there exists $p \in Q_0$ such that $(t_i \ldots t_1 \Phi^m(x))_p < 0$.

We assume that $p \leq i$. By Corollary 4.5, we have $\Phi = t_l \ldots t_1$. Then since t_j changes the only j-th coordinate, we have

$$(\Phi^{m+1}(x))_p = (t_l \ldots t_{i+1}(t_i \ldots t_1 \Phi^m(x)))_p < 0.$$

Similarly, if $p > i$, then we have $(\Phi^m(x))_p < 0$. Thus (1) implies (2). $\square$

Next we give a bijection between Φ_{Λ^i}-positive roots of q_{Λ^i} for $i \in Q_0$. In particular, this is given by reflection transformations. This is an analogue of a bijection between cluster-indecomposable modules of $\operatorname{mod} \Lambda^i$, which is given by reflection functors (Corollary 2.11). Thus next proposition implies that Φ-positive roots behave like cluster-indecomposable modules.

Proposition 5.10. *We have the following bijection for any $i \in Q_0$*

$$\{\Phi_{\Lambda^i}\text{-positive roots of } q_{\Lambda^i}\} \overset{1-1}{\longleftrightarrow} \{\Phi_{\Lambda^{i+1}}\text{-positive roots of } q_{\Lambda^{i+1}}\}$$

given by

$$x \longmapsto t_i x.$$

Proof. We only have to show for $i = 1$. Let x be a Φ_{Λ^1}-positive root of q_{Λ^1}. We will show that $t_1 x$ is a Φ_{Λ^2}-positive root of q_{Λ^2}. By $\Lambda^2 = \operatorname{End}_\Lambda(T_1)$ and Proposition 4.2, we have $\Phi_{\Lambda^2} = t_1 t_l \dots t_2$. Since x is a Φ_{Λ^1}-positive, we have

$$\Phi_{\Lambda^1}^m(x) = (t_l \dots t_1)^m(x) = t_l \dots t_2 \Phi_{\Lambda^2}^{m-1}(t_1 x) \in \mathbb{Z}_{\geq 0}^l$$

for any $m \in \mathbb{Z}$. Then by Lemma 3.9 and Lemma 5.9, $t_1 x$ is a Φ_{Λ^2}-positive root of q_{Λ^2}. Similarly, for a Φ_{Λ^2}-positive root y of q_{Λ^2}, we can show that $t_1^- y$ is a Φ_{Λ^1}-positive root. This implies the assertion. $\square$

As a corollary, we obtain the following result.

Corollary 5.11. *If Conjecture 5.3 is true for Λ, then it is also true for Λ^i for any $i \in Q_0$.*

Proof. We only have to show for $i = 2$. Let $y \in \mathbb{Z}^l$ be a Φ_{Λ^2}-positive root of q_{Λ^2}. It is enough to show that there exists a cluster-indecomposable Λ^2-module Y such that $\underline{\dim} Y = y$. By Proposition 5.10, there exists $x \in \mathbb{Z}^l$ such that $y = t_1 x$, where x is a Φ_{Λ^1}-positive root of q_{Λ^1}. On the other hand, there exists a cluster-indecomposable Λ-module X such that $\underline{\dim} X = x$ by the assumption. First, assume that $x \neq \mathbf{e}_1$, that is, $X \ncong P_1$. Then, by Corollary 2.11 and Corollary 3.8, we have a cluster-indecomposable Λ^2-module $\mathbf{T}_1^+ X$ such that $\underline{\dim}(\mathbf{T}_1^+ X) = t_1 x$. Next, assume that $x = \mathbf{e}_1$ and hence $t_1 x = \mathbf{e}_1 = y$. Since the vertex 1 is a source of the quiver of Λ^2, we have the simple injective module I_1 and $\underline{\dim} I_1 = y$. $\square$

Next, for the proof of Theorem 5.7, we prove the following key lemma.

Lemma 5.12. *Let $\Lambda = KQ/(R)$ be an n-representation-finite algebra and X be an indecomposable Λ-module. For any integer $u \geq 0$ and $i \in Q_0$, if $\mathbf{T}_{i-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^u(X)$ is either in $\mathcal{F}_0(T_i)$ or isomorphic to the simple projective Λ^i-module S_i, then X is cluster-indecomposable.*

Proof. First we will show that there exists an integer $v \geq 0$ and $j \in Q_0$ such that $\mathbf{T}_{j-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X)$ is isomorphic to the simple projective Λ^j-module S_j.

By Proposition 2.5 (5) and Proposition 4.7, we have $\mathbf{C}^{v+1}(X) = 0$ for a sufficiently large $v \geq 0$. Thus there exists an integer v and $j \in Q_0$ such that $\mathbf{T}_{j-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X) \neq 0$ and $\mathbf{T}_j^+ \mathbf{T}_{j-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X) = 0$. Hence we obtain $\mathbf{T}_{j-1}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X) \cong S_j$.

Then, because we have $\mathbf{T}_{j-2}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X) \in \mathcal{F}_0(T_{j-1})$, we get $\mathbf{T}_{j-1}^+ \mathbf{T}_{j-2}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X) \in \mathcal{X}_0(T_{j-1})$ by Lemma 2.9. Therefore we have

$$\mathbf{T}_{j-1}^- S_j \cong \mathbf{T}_{j-1}^- (\mathbf{T}_{j-1}^+ \mathbf{T}_{j-2}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X)) \cong \mathbf{T}_{j-2}^+ \dots \mathbf{T}_1^+ \mathbf{C}^v(X).$$

Applying this process repeatedly, we get $X \cong \mathbf{C}^{-v} \mathbf{T}_1^- \dots \mathbf{T}_{j-1}^-(S_j)$. Hence by Corollary 2.11, X is a cluster-indecomposable Λ-module. $\square$

Next we introduce the following notion.

Definition 5.13. Let Λ be a finite dimensional algebra. A *path* in $\operatorname{mod}\Lambda$ is a sequence

$$X_0 \xrightarrow{f_1} X_1 \to \cdots \to X_{c-1} \xrightarrow{f_c} X_c$$

of nonzero non-isomorphisms $f_1, \dots, f_c$ between indecomposable Λ-modules $X_0, \dots, X_c$. A path in $\operatorname{mod}\Lambda$ is called a *cycle* if $X_0 \cong X_c$. An algebra Λ is called *representation-directed* if there is no cycle in $\operatorname{mod}\Lambda$.

We recall an important theorem of representation-directed algebras.

Theorem 5.14 (Bongartz)**.** *Let Λ be a representation-directed algebra with $\operatorname{gl.dim}\Lambda \leq 2$ and q_Λ be the Euler form of Λ. The map $X \mapsto \underline{\dim} X$ gives a bijection between the isomorphism classes of indecomposable Λ-modules and the positive roots of q_Λ.*

Next lemma plays an important role to prove Theorem 5.7.

Lemma 5.15. *Let Λ be a 2-representation-finite algebra and $x \in \mathbb{Z}^l$ be a Φ-positive root of q_Λ. Assume that Λ is representation-directed. Then there exists an indecomposable Λ-module X such that $\underline{\dim} X = x$. Moreover, X satisfies one of the following conditions.*

(1) $X \in \mathcal{F}_0(T_1)$.
(2) $X \cong P_1$.

Proof. By Theorem 5.14, there exists an indecomposable Λ-module X such that $\underline{\dim} X = x$. We assume that $X \not\cong P_1$ and show that $X \in \mathcal{F}_0(T_1)$.

First we have

$$\begin{aligned}(t_1(\underline{\dim} X))_1 &= -\dim Xe_1 + \langle \underline{\dim} X, \mathbf{e}_1\rangle + \langle \mathbf{e}_1, \underline{\dim} X\rangle \\ &= \langle \underline{\dim} X, \mathbf{e}_1\rangle \\ &= (-1)^2 \langle \Phi^-(\mathbf{e}_1), \underline{\dim} X\rangle \\ &= \langle \underline{\dim}(\tau_n^- P_1), \underline{\dim} X\rangle \\ &= \sum_{j \geq 0} (-1)^j \dim \operatorname{Ext}^j_\Lambda(\tau_n^- P_1, X).\end{aligned}$$

Then by Lemma 2.2, we obtain

$$(t_1 x)_1 = \dim \operatorname{Hom}_\Lambda(\tau_2^- P_1, X) - \dim \operatorname{Ext}^1_\Lambda(\tau_2^- P_1, X) + \dim \operatorname{Ext}^2_\Lambda(\tau_2^- P_1, X).$$

By Proposition 2.5 (3), we obtain $\operatorname{Ext}^2_\Lambda(\tau_2^- P_1, X) \cong D\overline{\operatorname{Hom}}_\Lambda(X, P_1)$. Therefore we have $\dim \operatorname{Ext}^2_\Lambda(\tau_2^- P_1, X) = 0$ since P_1 is a simple projective Λ-module.

Now we assume that $X \notin \mathcal{F}_0(T_1)$. Then we have $\dim \operatorname{Ext}^1_\Lambda(\tau_2^- P_1, X) \neq 0$. Because we have $\operatorname{Ext}^1_\Lambda(\tau_2^- P_1, X) \cong D\overline{\operatorname{Hom}}_\Lambda(X, \tau(\tau_2^- P_1))$, there exists a path from X to $\tau(\tau_2^- P_1)$.

On the other hand, $\tau_2^- P_1$ is an indecomposable module, which is not projective by Proposition 2.5. Therefore we have an almost split sequence

$$0 \to \tau(\tau_2^- P_1) \to * \to \tau_2^- P_1 \to 0.$$

Hence there exists a path from $\tau(\tau_2^- P_1)$ to $\tau_2^- P_1$.

Then since Λ is representation-directed, there is no path from $\tau_2^- P_1$ to X and we get $\dim \operatorname{Hom}_\Lambda(\tau_2^- P_1, X) = 0$. Consequently we have $(t_1 x)_1 < 0$, which contradicts the fact that x is Φ-positive by Lemma 5.9. □

Then we give the proof of Theorem 5.7.

Proof. Let $x \in \mathbb{Z}^l$ be a Φ-positive root of q_Λ. By Theorem 5.14, there exists an indecomposable Λ-module X such that $\underline{\dim} X = x$. We will show that X is a cluster-indecomposable module. It is enough to show that X satisfies the condition of Lemma 5.12.

If $X \cong P_1$, then there is nothing to show. If $X \ncong P_1$, then we have $X \in \mathcal{F}_0(T_1)$ by Lemma 5.15. Then by Proposition 3.7, $\mathbf{T}_1^+ X$ is an indecomposable Λ^2-module and we have $\underline{\dim}(\mathbf{T}_1^+ X) = t_1 x$. By Proposition 5.10, $t_1 x$ is a Φ_{Λ^2}-positive root of q_{Λ^2} and we can apply Lemma 5.15 again. Thus X satisfies the condition of Lemma 5.12 and hence X is a cluster-indecomposable Λ-module. $\square$

Then Corollary 5.8 follows immediately from this result.

Proposition 5.16. *Let Λ be a 2-representation-finite algebra. If Λ is iterated tilted, then Λ^i is representation-directed for any $i \in Q_0$.*

Proof. By [IO2, Theorem 3.12] (see also [Rin2]), Λ is derived equivalent to a path algebra of a Dynkin quiver. Therefore Λ and hence Λ^i are representation-directed for any $i \in Q_0$. $\square$

Example 5.17. (1) Let $\Lambda = KQ/(R)$ be the 2-representation-finite algebra given by the following quiver with relations.

$Q =$



$$R = \{c_1c_2 - b_5b_4b_3b_2b_1, b_ia_i\}.$$

This algebra is iterated tilted by [IO2, Theorem 3.12].

Then there are 57 indecomposable Λ-modules, and their dimension vectors give the following complete list of positive roots of q_Λ.

$$\begin{pmatrix}00000\\100000\\0\end{pmatrix}, \begin{pmatrix}00000\\010000\\0\end{pmatrix}, \begin{pmatrix}00000\\110000\\0\end{pmatrix}, \begin{pmatrix}00000\\001000\\0\end{pmatrix}, \begin{pmatrix}00000\\011000\\0\end{pmatrix}, \begin{pmatrix}00000\\111000\\0\end{pmatrix}, \begin{pmatrix}00000\\000100\\0\end{pmatrix}, \begin{pmatrix}00000\\001100\\0\end{pmatrix},$$
$$\begin{pmatrix}00000\\011100\\0\end{pmatrix}, \begin{pmatrix}00000\\111100\\0\end{pmatrix}, \begin{pmatrix}00000\\000010\\0\end{pmatrix}, \begin{pmatrix}00000\\000110\\0\end{pmatrix}, \begin{pmatrix}00000\\001110\\0\end{pmatrix}, \begin{pmatrix}00000\\011110\\0\end{pmatrix}, \begin{pmatrix}00000\\111110\\0\end{pmatrix}, \begin{pmatrix}00000\\000000\\1\end{pmatrix},$$
$$\begin{pmatrix}00000\\100000\\1\end{pmatrix}, \begin{pmatrix}00000\\110000\\1\end{pmatrix}, \begin{pmatrix}00000\\111000\\1\end{pmatrix}, \begin{pmatrix}00000\\111100\\1\end{pmatrix}, \begin{pmatrix}00000\\111110\\1\end{pmatrix}, \begin{pmatrix}00000\\000001\\0\end{pmatrix}, \begin{pmatrix}00000\\000011\\0\end{pmatrix}, \begin{pmatrix}00000\\000111\\0\end{pmatrix},$$
$$\begin{pmatrix}00000\\001111\\0\end{pmatrix}, \begin{pmatrix}00000\\011111\\0\end{pmatrix}, \begin{pmatrix}00000\\000001\\1\end{pmatrix}, \begin{pmatrix}00000\\000011\\1\end{pmatrix}, \begin{pmatrix}00000\\000111\\1\end{pmatrix}, \begin{pmatrix}00000\\001111\\1\end{pmatrix}, \begin{pmatrix}00000\\011111\\1\end{pmatrix}, \begin{pmatrix}00000\\111111\\1\end{pmatrix},$$
$$\begin{pmatrix}10000\\000000\\0\end{pmatrix}, \begin{pmatrix}10000\\010000\\0\end{pmatrix}, \begin{pmatrix}10000\\011000\\0\end{pmatrix}, \begin{pmatrix}10000\\011100\\0\end{pmatrix}, \begin{pmatrix}10000\\011110\\0\end{pmatrix}, \begin{pmatrix}10000\\011111\\0\end{pmatrix}, \begin{pmatrix}10000\\011111\\1\end{pmatrix}, \begin{pmatrix}01000\\000000\\0\end{pmatrix},$$
$$\begin{pmatrix}01000\\001000\\0\end{pmatrix}, \begin{pmatrix}01000\\001100\\0\end{pmatrix}, \begin{pmatrix}01000\\001110\\0\end{pmatrix}, \begin{pmatrix}01000\\001111\\0\end{pmatrix}, \begin{pmatrix}01000\\001111\\1\end{pmatrix}, \begin{pmatrix}00100\\000000\\0\end{pmatrix}, \begin{pmatrix}00100\\000100\\0\end{pmatrix}, \begin{pmatrix}00100\\000110\\0\end{pmatrix},$$
$$\begin{pmatrix}00100\\000111\\0\end{pmatrix}, \begin{pmatrix}00100\\000111\\1\end{pmatrix}, \begin{pmatrix}00010\\000000\\0\end{pmatrix}, \begin{pmatrix}00010\\000010\\0\end{pmatrix}, \begin{pmatrix}00010\\000011\\0\end{pmatrix}, \begin{pmatrix}00010\\000011\\1\end{pmatrix}, \begin{pmatrix}00001\\000000\\0\end{pmatrix}, \begin{pmatrix}00001\\000001\\0\end{pmatrix},$$
$$\begin{pmatrix}00001\\000001\\1\end{pmatrix}.$$

Moreover there are 22 cluster-indecomposable Λ-modules, and their dimension vectors give the following complete list of Φ-positive roots.

$$\begin{pmatrix}00001\\000000\\0\end{pmatrix}, \begin{pmatrix}00010\\000000\\0\end{pmatrix}, \begin{pmatrix}00100\\000000\\0\end{pmatrix}, \begin{pmatrix}01000\\000000\\0\end{pmatrix}, \begin{pmatrix}10000\\000000\\0\end{pmatrix}, \begin{pmatrix}00001\\000001\\0\end{pmatrix}, \begin{pmatrix}00001\\000001\\1\end{pmatrix}, \begin{pmatrix}00010\\000011\\0\end{pmatrix},$$
$$\begin{pmatrix}00100\\000111\\0\end{pmatrix}, \begin{pmatrix}01000\\001111\\0\end{pmatrix}, \begin{pmatrix}10000\\011111\\0\end{pmatrix}, \begin{pmatrix}00000\\111111\\1\end{pmatrix}, \begin{pmatrix}00010\\000010\\0\end{pmatrix}, \begin{pmatrix}00100\\000100\\0\end{pmatrix}, \begin{pmatrix}01000\\001000\\0\end{pmatrix}, \begin{pmatrix}10000\\010000\\0\end{pmatrix},$$
$$\begin{pmatrix}00000\\100000\\1\end{pmatrix}, \begin{pmatrix}00000\\111110\\0\end{pmatrix}, \begin{pmatrix}00000\\111100\\0\end{pmatrix}, \begin{pmatrix}00000\\111000\\0\end{pmatrix}, \begin{pmatrix}00000\\110000\\0\end{pmatrix}, \begin{pmatrix}00000\\100000\\0\end{pmatrix}.$$

(2) Let Λ be the 2-representation-finite algebra given as an Auslander algebra of A_6.



In this case, there exist infinitely many indecomposable Λ-modules and infinitely many roots of q_Λ. However, by Theorem 4.10, one can see that there are 56 cluster-roots. On the other hand, one can see that there are 56 Φ-positive roots, for example, by Mathematica and they correspond bijectively to the cluster-roots.

6. Quivers of n-APR tilts and reflection functors

One of the important features of APR tilting modules over a path algebra is that we can describe the quiver of its endomorphism algebra. More precisely, the new quiver is obtained by reversing the arrows adjacent to the sink of the original quiver [BGP, APR]. In this section, we provide a generalization of this result to n-APR tilting modules and give an explicit description of n-APR tilts by quivers with relations. Moreover, using this, we introduce reflection functors by linear representations of quivers.

Throughout this section, let $\Lambda = KQ/(R)$ be a finite dimensional algebra with a minimal set R of relations. Let $k \in Q_0$ be a sink admitting the n-APR tilting module T_k. Take a minimal injective resolution of P_k

$$\begin{tikzcd} 0 \arrow[r] & P_k \arrow[r] & I^0 \arrow[r] & \dots \arrow[r] & I^{n-1} \arrow[r] & I^n \arrow[r] & 0. \end{tikzcd}$$

Then, by [IO1, Theorem 3.2], we have a minimal projective resolution of $\tau_n^- P_k$

(i)
$$\begin{tikzcd} 0 \arrow[r] & P^0 \arrow[r] & \dots \arrow[r] & P^{n-1} \arrow[r, "\alpha"] & P^n \arrow[r, "\beta"] & \tau_n^- P_k \arrow[r] & 0, \end{tikzcd}$$

where $P^i = \nu^-(I^i)$ for $0 \le i \le n$. Note that we have $P^i \in \mathrm{add}(\Lambda/P_k)$ for $1 \le i \le n$ since P_k is a simple projective Λ-module.

We decompose P^{n-1} and P^n into direct sums of indecomposable modules and denote them by $\bigoplus_{b\in B} P_{i_b}$ and $\bigoplus_{c\in C} P_{i_c}$, respectively. We denote the morphism α by $(a_{cb})_{cb} : \bigoplus_{b\in B} P_{i_b} \to \bigoplus_{c\in C} P_{i_c}$ and we define new arrows $a_c^* : k \to i_c$ for each $c \in C$. Then we give the following definition.

Definition 6.1. Under the above set-up, we define a new quiver with relations $(Q', R') = \sigma_k(Q, R)$ as follows.

(1) $Q'_0 = Q_0$
(2) $Q'_1 = \{a \in Q_1 \mid e(a) \ne k\} \coprod \{\, a_c^* : k \to i_c \mid c \in C\}$.
(3) $R' = \begin{cases} 0 & \text{if } n = 1 \\ \{r \in R \mid e(r) \ne k\} \coprod \{\, \sum_{c\in C} a_c^* a_{cb} \mid b \in B\} & \text{if } n > 1. \end{cases}$

Similarly, for a source k admitting the n-APR co-tilting module, we define σ_k^-.

Note that if $n = 1$, Q' is obtained just reversing the arrows and if $n = 2$, (Q', R') is determined by a simple method (cf. [IO1, section 3.3]).

Example 6.2. (1) Let $\Lambda = KQ/(R)$ be the algebra given by the following quiver with relations

$$\begin{tikzcd}
 & & 3 \arrow[dl, "a_1"'] \\
1 & 2 \arrow[l, "a_3"'] & \\
 & & 4, \arrow[ul, "a_2"]
\end{tikzcd} \qquad R = \{a_1a_3, a_2a_3\}.$$

Then P_1 admits a 2-APR tilting module and we have the following projective resolution

$$\begin{tikzcd}
0 \arrow[r] & P_1 \arrow[r] & P_2 \arrow[r] & P_3 \oplus P_4 \arrow[r] & \tau_2^- P_1 \arrow[r] & 0.
\end{tikzcd}$$

Then $(Q', R') = \sigma_1(Q, R)$ is given as follows

$$\begin{tikzcd}
 & & 3 \arrow[dl, "a_1"] \\
1 \arrow[urr, bend left, "b_1"] \arrow[drr, bend right, "b_2"'] & 2 & \\
 & & 4, \arrow[ul, "a_2"']
\end{tikzcd} \qquad R' = \{b_1a_1, b_2a_2\}.$$

(2) Let $\Lambda = KQ/(R)$ be the algebra given by the following quiver with relations

$$\begin{tikzcd}
 & & 3 \arrow[dl, "b_1"'] & \\
1 & 2 \arrow[l, "a"] & & 5 \arrow[ul, "c_1"'] \arrow[dl, "c_2"] \\
 & & 4, \arrow[ul, "b_2"] &
\end{tikzcd} \qquad R = \{b_1a, b_2a, c_1b_1 - c_2b_2\}.$$

Then P_1 admits a 3-APR tilting module and we have the following projective resolution

$$\begin{tikzcd}
0 \arrow[r] & P_1 \arrow[r] & P_2 \arrow[r] & P_3 \oplus P_4 \arrow[r] & P_5 \arrow[r] & \tau_3^- P_1 \arrow[r] & 0.
\end{tikzcd}$$

Then $(Q', R') = \sigma_1(Q, R)$ is given as follows

$$\begin{tikzcd}
 & & 3 \arrow[dl, "b_1"'] & \\
1 \arrow[rrr, bend left, "d"] & 2 & & 5 \arrow[ul, "c_1"'] \arrow[dl, "c_2"] \\
 & & 4, \arrow[ul, "b_2"] &
\end{tikzcd} \qquad R' = \{dc_1, dc_2, c_1b_1 - c_2b_2\}.$$

Then we give the following proposition.

Proposition 6.3. *Let $\Lambda = KQ/(R)$ be a finite dimensional algebra with a minimal set R of relations. Assume that $k \in Q_0$ is a sink admitting an n-APR tilting module T_k and $(Q', R') := \sigma_k(Q, R)$. Then there is an isomorphism $\mathrm{End}_\Lambda(T_k) \cong KQ'/(R')$ and R' is a minimal set of relations.*

In this proposition, if $KQ/(R)$ is n-representation-finite, then so is $KQ'/(R')$ by Proposition 2.8 (see Example 2.13). Note that the proof was given by [IO1, Theorem 3.11] for $n = 2$. Our proof is quite different. Let us start with recalling the following result. We denote by J_Λ the Jacobson radical of an algebra Λ and call an element of KQ *basic* if it is a linear sum of paths in Q with a common start and a common end.

Theorem 6.4. [BIRS, Proposition 3.1, 3.3] *Let Q be a finite connected acyclic quiver and Γ be a basic finite dimensional algebra. Let $\phi : KQ \to \Gamma$ be an algebra homomorphism and R be a finite set of basic elements in J_{KQ}. Then the following conditions are equivalent.*

(1) *ϕ is surjective and* $\mathrm{Ker}\,\phi = I$ *for the ideal $I = (R)$ of KQ.*

(2) *The following sequence is exact for any $i \in Q_0$*

$$\bigoplus_{r\in R, s(r)=i} (\phi e(r))\Gamma \xrightarrow{(\phi\partial_a r)} \bigoplus_{a\in Q_1, s(a)=i} (\phi e(a))\Gamma \xrightarrow{(\phi a)} (\phi i)J_\Gamma \longrightarrow 0.$$

Then we give a proof of Proposition 6.3.

Proof. Take the minimal projective resolution (i) and follow the previous notations. We write $\Gamma := \operatorname{End}_\Lambda(T_k)$. We denote by ϕ the natural surjective map $KQ \to \Lambda$, which gives $\operatorname{Ker}\phi = (R)$. Then we define an algebra homomorphism $\phi' : KQ' \to \Gamma$ as follows.

(a) For $i \in Q'_0$, denote the canonical projection by $\pi_i : T_k \to P_i$ for $i \neq k$ and $\pi_k : T_k \to \tau_n^- P_k$, and denote the canonical injection by $\iota_i : P_i \to T_k$ for $i \neq k$ and $\iota_k : \tau_n^- P_k \to T_k$. Then, we define $\phi' i = \iota_i \pi_i \in \Gamma$, which gives an idempotent of Γ.

(b) For $a \in Q'_1 \cap Q_1$, define $\phi' a = \phi a$. For a^*_c with $c \in C$, define

$$\phi'((a^*_c)_{c\in C}) = \beta \in \operatorname{Hom}_\Lambda(\bigoplus_{c\in C} P_{i_c}, \tau_n^- P_k),$$

where β is given in (i).

Take a vertex $i \in Q_0$ with $i \neq k$ and let S_i be the simple Λ-module. Then we have a minimal projective presentation of S_i as follows

$$\text{(ii)} \qquad \bigoplus_{r\in R, s(r)=i} P_{e(r)} \xrightarrow{f_i} \bigoplus_{a\in Q_1, s(a)=i} P_{e(a)} \xrightarrow{g_i} P_i \longrightarrow S_i \longrightarrow 0.$$

Then by Proposition [IO1, Theorem 3.16], we have a projective presentation of a simple Γ-module S^Γ_i
(iii)

$$\operatorname{Hom}_\Lambda(T_k, \bigoplus_{r\in R, s(r)=i} P_{e(r)}) \xrightarrow{\cdot f_i} \operatorname{Hom}_\Lambda(T_k, \bigoplus_{a\in Q_1, s(a)=i} P_{e(a)}) \xrightarrow{\cdot g_i} \operatorname{Hom}_\Lambda(T_k, P_i) \longrightarrow S^\Gamma_i \longrightarrow 0,$$

and of a simple Γ-module S^Γ_k
(iv)

$$\operatorname{Hom}_\Lambda(T_k, \textstyle\bigoplus_{b\in B} P_{i_b}) \xrightarrow{\cdot\alpha} \operatorname{Hom}_\Lambda(T_k, \bigoplus_{c\in C} P_{i_c}) \xrightarrow{\cdot\beta} \operatorname{Hom}_\Lambda(T_k, \tau_n^- P_k) \longrightarrow S^\Gamma_k \longrightarrow 0.$$

Note that we have $\operatorname{Hom}_\Lambda(T_k, P_k) = 0$ since P_k is a simple projective module. Then by Theorem 6.4, the first statement follows. Moreover, (iii),(iv) are minimal projective presentations. Indeed, we have an equivalence $\operatorname{add}(T_k) \cong \operatorname{proj}(\Gamma)$, where $\operatorname{proj}(\Gamma)$ denote the category of projective modules of $\operatorname{mod}\Gamma$. Then if (iii),(iv) are not minimal, it implies that the sequence (i),(ii) are not minimal, which is a contradiction. Then by [BIRS], the second statement follows. □

Next, following [BGP], we define reflection functors in terms of linear representations. First we recall basic results. Let $\Lambda = KQ/(R)$ be a finite dimensional algebra and denote by $\operatorname{rep}_K(Q, R)$ the category of finite dimensional K-linear representations of Q bound by R (refer to [ASS] for the precise definition). Then we recall the following fundamental theorem.

Theorem 6.5. *Let $\Lambda = KQ/(R)$ be a finite dimensional algebra. There exists a K-linear equivalence of categories*

$$F : \operatorname{mod}\Lambda \to \operatorname{rep}_K(Q, R).$$

Recall that the functor F associates with any Λ-module M the representation $FM = ((FM)_i, \varphi_a)_{i\in Q_0, a\in Q_1}$, where $(FM)_i = Me_i$ for $i \in Q_0$ and the K-linear map $\varphi_a : Me_i \to Me_j$ for $a : i \to j$ in Q_1 is defined by $x \mapsto xa$ for $x \in Me_i$.

Definition 6.6. (I) Let $\Lambda = KQ/(R)$ be a finite dimensional algebra, $k \in Q_0$ be a sink admitting the n-APR tilting module T_k and $(Q', R') := \sigma_k(Q, R)$. We define the *reflection functor*

$$\mathbf{Q}_k^+ : \mathrm{rep}_K(Q, R) \to \mathrm{rep}_K(Q', R')$$

as follows.

Take the minimal projective resolution (i) and follow the previous notations. For an object $X = (X_i, \varphi_a)$ in $\mathrm{rep}_K(Q, R)$, we define the object $\mathbf{Q}_k^+(X) = (X_i', \varphi_a')$ in $\mathrm{rep}_K(Q', R')$ as follows.

(1) $X_i' = X_i$ for $i \neq k$, whereas X_k' is the kernel of the K-linear map

$$(\varphi_{a_{cb}})_{a_{cb}} : \bigoplus_{c\in C} X_{i_c} \to \bigoplus_{b\in B} X_{i_b},$$

where $\varphi_{a_{cb}}$ is defined by φ_a for $a \in Q_1$ naturally .

(2) $\varphi_a' = \varphi_a$ for all arrows $a : i \to j$ in Q_1 with $j \neq k$, whereas, for $a_c^* : k \to i_c$ in Q_1', define $\varphi_{a_c^*} : X_k' \to X_{i_c}$ by the composition of the inclusion of X_k' into $\bigoplus_{c\in C} X_{i_c}$ with the projection onto the direct summand X_{i_c}.

Next let $f = (f_i)_{i\in Q_0} : X \to Y$ be a morphism in $\mathrm{rep}_K(Q, R)$, where $X = (X_i, \varphi_a)$ and $Y = (Y_i, \psi_a)$. We define the morphism

$$\mathbf{Q}_k^+(f) = f' = (f_i')_{i\in Q_0'} : \mathbf{Q}_k^+(X) \to \mathbf{Q}_k^+(Y)$$

in $\mathrm{rep}_K(Q', R')$ as follows. For all $i \neq k$, we let $f_i' = f_i$, whereas f_k' is the unique K-linear map, making the following diagram commutative

$$\begin{tikzcd}
0 \arrow[r] & (\mathbf{Q}_k^+(X))_k \arrow[r] \arrow[d, dotted, "f_k'"] & \bigoplus_{c\in C} X_{i_c} \arrow[r, "(\varphi_{a_{cb}})_{a_{cb}}"] \arrow[d, "\oplus f_c"] & \bigoplus_{b\in B} X_{i_b} \arrow[d, "\oplus f_b"] \\
0 \arrow[r] & (\mathbf{Q}_k^+(Y))_k \arrow[r] & \bigoplus_{c\in C} Y_{i_c} \arrow[r, "(\psi_{a_{cb}})_{a_{cb}}"] & \bigoplus_{b\in B} Y_{i_b}.
\end{tikzcd}$$

(II) Let $\Gamma = KQ'/(R')$ be a finite dimensional algebra, $k \in Q_0'$ be a source admitting the n-APR co-tilting module T_k and $(Q, R) := \sigma_k^-(Q', R')$. We define the *reflection functor*

$$\mathbf{Q}_k^- : \mathrm{rep}_K(Q', R') \to \mathrm{rep}_K(Q, R)$$

as follows.

We have a minimal injective resolution of $\tau_n I_k$

$$0 \longrightarrow \tau_n I_k \longrightarrow I^n \xrightarrow{\gamma} I^{n-1} \longrightarrow \dots \longrightarrow I^0 \longrightarrow 0.$$

We decompose I^n and I^{n-1} into direct sums of indecomposable modules, and denote them by $\bigoplus_{c\in C} I_{i_c}$ and $\bigoplus_{b\in B} I_{i_b}$, respectively. We denote the morphism γ by $(\varphi_{a_{bc}})_{a_{bc}} : \bigoplus_{c\in C} I_{i_c} \to \bigoplus_{b\in B} I_{i_b}$. For an object $X' = (X_i', \varphi_a')$ in $\mathrm{rep}_K(Q', R')$, we define the object $\mathbf{Q}_k^-(X') = (X_i, \varphi_a)$ in $\mathrm{rep}_K(Q, R)$ as follows.

(1) $X_i = X'_i$ for $i \neq k$, whereas X_k is the cokernel of the K-linear map

$$(\varphi_{a_{bc}})_{a_{bc}} : \bigoplus_{b\in B} X'_{i_b} \to \bigoplus_{c\in C} X'_{i_c}.$$

(2) $\varphi_a = \varphi'_a$ for all arrows $a : i \to j$ in Q' with $i \neq k$, whereas, for $a^*_c : i_c \to k$ in Q_1, define $\varphi_{a^*_c} : X'_{i_c} \to X_k$ by the composition of the inclusion of X'_{i_c} into $\bigoplus_{c\in C} X'_{i_c}$ with the cokernel projection onto X_k.

Next let $f' = (f'_i)_{i\in Q'_0} : X' \to Y'$ be a morphism in $\mathrm{rep}_K(Q', R')$, where $X' = (X'_i, \varphi'_a)$ and $Y' = (Y'_i, \psi'_a)$. We define the morphism

$$\mathbf{Q}^-_k(f') = f = (f_i)_{i\in Q_0} : \mathbf{Q}^-_k(X') \to \mathbf{Q}^-_k(Y')$$

in $\mathrm{rep}_K(Q, R)$ as follows. For all $i \neq k$, we let $f_i = f'_i$, whereas f_k is the unique K-linear map, making the following diagram commutative

$$\begin{tikzcd}
\bigoplus_{b\in B} X'_{i_b} \arrow[r, "(\varphi_{a_{bc}})_{a_{bc}}"] \arrow[d, "\oplus f_b"] & \bigoplus_{c\in C} X'_{i_c} \arrow[r] \arrow[d, "\oplus f_c"] & (\mathbf{Q}^-_k(X'))_k \arrow[r] \arrow[d, dotted, "f_k"] & 0 \\
\bigoplus_{b\in B} Y'_{i_b} \arrow[r, "(\varphi_{a_{bc}})_{a_{bc}}"] & \bigoplus_{c\in C} Y'_{i_c} \arrow[r] & (\mathbf{Q}^-_k(Y'))_k \arrow[r] & 0.
\end{tikzcd}$$

Example 6.7. Let $\Lambda = KQ/(R)$ be the algebra given by Example 6.2 (2) and X be a representation of $\mathrm{rep}_K(Q, R)$ given as follows

$$\begin{tikzcd}
 & & X_3 \arrow[dl, "\varphi_{b_1}"'] & \\
X_1 & X_2 \arrow[l, "\varphi_a"] & & X_5 \arrow[ul, "\varphi_{c_1}"'] \arrow[dl, "\varphi_{c_2}"] \\
 & & X_4 \arrow[ul, "\varphi_{b_2}"] &
\end{tikzcd}$$

Then $\mathbf{Q}^+_k(X)$ is given as follows

$$\begin{tikzcd}
 & & X_3 \arrow[dl, "\varphi_{b_1}"'] & \\
X'_1 \arrow[rrr, bend left, "\varphi_d"] & X_2 & & X_5 \arrow[ul, "\varphi_{c_1}"'] \arrow[dl, "\varphi_{c_2}"] \\
 & & X_4 \arrow[ul, "\varphi_{b_2}"] &
\end{tikzcd}$$

where $X'_1 = \mathrm{Ker}\, F$ for $F : X_5 \xrightarrow{\binom{\varphi_{c_1}}{\varphi_{c_2}}} X_3 \oplus X_4$.

Finally we give a relationship between $\mathbf{Q}^+_k, \mathbf{Q}^-_k$ and $\mathbf{T}^+_k := \mathrm{Hom}_\Lambda(T_k, -), \mathbf{T}^-_i := - \otimes_\Gamma T_k$ as follows.

Proposition 6.8. *Let $\Lambda = KQ/(R)$ be a finite dimensional algebra and $k \in Q_0$ be a sink admitting an n-APR tilting module T_k. We denote by $(Q', R') := \sigma_k(Q, R)$ and $\Gamma := KQ'/(R')$. Then we have $\mathbf{Q}^+_k \circ F \cong F' \circ \mathbf{T}^+_k$ on $\mathcal{F}_0(T_k)$ and $\mathbf{Q}^-_k \circ F' \cong F \circ \mathbf{T}^-_k$ on*

$\mathcal{X}_0(T_k)$, where F and F' are the category equivalences.

$$\begin{tikzcd}
\operatorname{mod} KQ/(R) \arrow[r, "\mathbf{T}_k^+"] \arrow[d, "F"', "\cong"] & \operatorname{mod} KQ'/(R') \arrow[r, "\mathbf{T}_k^-"] \arrow[d, "F'"', "\cong"] & \operatorname{mod} KQ/(R) \arrow[d, "F"', "\cong"] \\
\operatorname{rep}_K(Q,R) \arrow[r, "\mathbf{Q}_k^+"] & \operatorname{rep}_K(Q',R') \arrow[r, "\mathbf{Q}_k^-"] & \operatorname{rep}_K(Q,R).
\end{tikzcd}$$

Proof. We only prove that $\mathbf{Q}_k^+ \circ F \cong F' \circ \mathbf{T}_k^+$; the proof of the second statement is similar. Let X be a Λ-module with $X \in \mathcal{F}_0(T_k)$. Take a vertex $i \neq k$. Then we have

$$\begin{aligned}
(F' \operatorname{Hom}_\Lambda(T_k, X))_i &\cong (\operatorname{Hom}_\Lambda(T_k, X))e_i \\
&\cong \operatorname{Hom}_\Gamma(e_i\Gamma, \operatorname{Hom}_\Lambda(T_k, X)) \\
&\cong \operatorname{Hom}_\Gamma(\operatorname{Hom}_\Lambda(T_k, e_i\Lambda), \operatorname{Hom}_\Lambda(T_k, X)) \\
&\cong \operatorname{Hom}_\Lambda(e_i\Lambda, X) \qquad \text{(Lemma 2.9)} \\
&\cong Xe_i \\
&\cong (FX)_i \\
&\cong (\mathbf{Q}_k^+(FX))_i.
\end{aligned}$$

On the other hand, if $i = k$, we have

$$\begin{aligned}
(F' \operatorname{Hom}_\Lambda(T_k, X))_k &\cong (\operatorname{Hom}_\Lambda(T_k, X))e_k \\
&\cong \operatorname{Hom}_\Gamma(e_k\Gamma, \operatorname{Hom}_\Lambda(T_k, X)) \\
&\cong \operatorname{Hom}_\Gamma(\operatorname{Hom}_\Lambda(T_k, \tau_n^- P_k), \operatorname{Hom}_\Lambda(T_k, X)) \\
&\cong \operatorname{Hom}_\Lambda(\tau_n^- P_k, X). \qquad \text{(Lemma 2.9)}
\end{aligned}$$

By applying $\operatorname{Hom}_\Lambda(-, X)$ to the minimal projective resolution (i), we have following commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \operatorname{Hom}_\Lambda(\tau_n^- P_k, X) \arrow[r] \arrow[d, dotted] & \operatorname{Hom}_\Lambda(\bigoplus_{c\in C} P_{i_c}, X) \arrow[r, "(a_{cb})_{a_{cb}}"] \arrow[d, "\cong"] & \operatorname{Hom}_\Lambda(\bigoplus_{b\in B} P_{i_b}, X) \arrow[d, "\cong"] \\
0 \arrow[r] & (\mathbf{Q}_k^+(FX))_k \arrow[r] & \bigoplus_{c\in C} (FX)_{i_c} \arrow[r, "(\varphi_{a_{cb}})_{a_{cb}}"] & \bigoplus_{b\in B} (FX)_{i_b}
\end{tikzcd}$$

Hence we have $(\mathbf{Q}_k^+(FX))_k \cong (F' \operatorname{Hom}_\Lambda(T_k, X))_k$ as a K-vector space. Then one can easily check that it induces an isomorphism of representations $\mathbf{Q}_k^+ F(X) \cong F'\mathbf{T}_k^+(X)$ and it is easy to verify that the isomorphism is functorial. Thus we have $\mathbf{Q}_k^+ \circ F \cong F' \circ \mathbf{T}_k^+$. $\square$

References

[ASS] I. Assem, D. Simson, A. Skowroński, *Elements of the Representation Theory of Associative Algebras. Vol. 1*, London Mathematical Society Student Texts 65, Cambridge university press (2006).

[APR] M. Auslander, M. I. Platzeck, I. Reiten, *Coxeter functors without diagrams*, Trans. Amer. Math. Soc. 250 (1979), 1–46.

[ARS] M. Auslander, I. Reiten, S. O. Smalø, *Representation Theory of Artin Algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1995.

[BGP] I. N. Bernstein, I. M. Gelfand, V. A. Ponomarev, *Coxeter functors and Gabriel's theorem*, Uspehi Mat. Nauk 28 (1973), no. 2(170), 19–33.

[B1] K. Bongartz, *Tilted algebras*, Representations of algebras (Puebla, 1980), 26–38, Lecture Notes in Math., 903, Springer, Berlin-New York, 1981.

[B2] K. Bongartz, *Algebras and quadratic forms*, J. London Math. Soc. (2) 28 (1983), no. 3, 461–469.

[B3] K. Bongartz, *A criterion for finite representation type*, Math. Ann. 269 (1984), no. 1, 1–12.

[BIRS] A. B. Buan, O. Iyama, I. Reiten, D. Smith, *Mutation of cluster-tilting objects and potentials*, Amer. J. Math. 133 (2011), no. 4, 835–887.

[BM] A. Buan, R. Marsh, *Cluster tilting theory*, Trends in representation theory of algebras and related topics, 1–30, Contemp. Math., 406, Amer. Math. Soc., Providence, RI, 2006.

[BB] S. Brenner, M. C. R. Butler, *Generalisations of the Bernstein-Gelfand-Ponomarev reflection functors*, in Proc. ICRA II (Ottawa,1979), Lecture Notes in Math. No. 832, Springer-Verlag, Berlin, Heidelberg, New York, 1980, pp. 103–69.

[DR] V. Dlab, C. M. Ringel, *Indecomposable representations of graphs and algebras*, Mem. Amer. Math. Soc. 6 (1976), no. 173.

[DF] P. Donovan, M. R. Freislich, *The representation theory of finite graphs and associated algebras*, Carleton Mathematical Lecture Notes, No. 5. Carleton University, Ottawa, Ont., 1973.

[D] P. Dräxler, *On indecomposable modules over directed algebras*, Proc. Amer. Math. Soc. 112 (1991), no. 2, 321–327.

[EH] K. Erdmann, T. Holm, *Maximal n-orthogonal modules for selfinjective algebras*, Proc. Amer. Math. Soc. 136 (2008) no. 9, 3069–3078.

[G] P. Gabriel, *Unzerlegbare Darstellungen I*, Manu. Math. 6 (1972), 71–103.

[H] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988.

[HR1] D. Happel, C. M. Ringel, *Tilted algebras*, Trans. Amer. Math. Soc. 274 (1982), no. 2, 399–443.

[HR2] D. Happel, C. M. Ringel, *Directing projective modules*, Arch. Math. (Basel) 60 (1993), no. 3, 237–246.

[HI1] M. Herschend, O. Iyama, *n-representation-finite algebras and twisted fractionally Calabi-Yau algebras*, Bull. Lond. Math. Soc. 43 (2011), no. 3, 449–466.

[HI2] M. Herschend, O. Iyama, *Selfinjective quivers with potential and 2-representation-finite algebras*, Compos. Math. 147 (2011), no. 6, 1885–1920.

[HIO] M. Herschend, O. Iyama, S. Oppermann, *n-representation infinite algebras*, arXiv:1205.1272.

[HZ] Z. Huang, X. Zhang, *The existence of maximal n-orthogonal subcategories*, J. Algebra 321 (2009), no. 10, 2829–2842.

[I1] O. Iyama, *Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories*, Adv. Math. 210 (2007), no. 1, 22–50.

[I2] O. Iyama, *Auslander correspondence*, Adv. Math. 210 (2007), no. 1, 51–82.

[I3] O. Iyama, *Cluster tilting for higher Auslander algebras*, Adv. Math. 226 (2011), no. 1, 1–61.

[IO1] O. Iyama, S. Oppermann, *n-representation-finite algebras and n-APR tilting*, Trans. Amer. Math. Soc. 363 (2011), no. 12, 6575–6614.

[IO2] O. Iyama, S. Oppermann, *Stable categories of higher preprojective algebras*, arXiv:0912.3412.

[M] Y. Miyashita, *Tilting modules of finite projective dimension*, Math. Z. 193 (1986), no. 1, 113–146.

[N] L. A. Nazarova, *Representations of quivers of infinite type*, Izv. Akad. Nauk SSSR Ser. Mat. 37 (1973), 752–791.

[Ka] V. G. Kac, *Infinite root systems, representations of graphs and invariant theory*, Invent. Math. 56 (1980), no. 1, 57–92.

[Ke1] B. Keller, *Cluster algebras, quiver representations and triangulated categories*, Triangulated categories, 76–160, London Math. Soc. Lecture Note Ser., 375, Cambridge Univ. Press, Cambridge, 2010.

[Ke2] B. Keller, *On cluster theory and quantum dilogarithm identities*, Representations of Algebras and Related Topics, EMS Series of Congress Reports. pp. 85–116.

[Kr] H. Krause, *Representations of quivers via reflection functors*, arXiv:0804.1428.

[RS] N. Reading, D. Speyer, *Combinatorial frameworks for cluster algebras*, arXiv:1111.2652.

[Re] I. Reiten, *Cluster categories*, Proceedings of the International Congress of Mathematicians. Volume I, 558–594, Hindustan Book Agency, New Delhi, 2010.

[Ric] J. Rickard, *Morita theory for derived categories*, J. London Math. Soc. (2) 39 (1989), no. 3, 436–456.

[Rin1] C. M. Ringel, *Tame algebras and integral quadratic forms*, Lecture Notes in Mathematics, 1099. Springer-Verlag, Berlin, 1984.

[Rin2] C. M. Ringel, *The self-injective cluster-tilted algebras*, Arch. Math. (Basel) 91 (2008), no. 3, 218–225.

Graduate School of Mathematics, Nagoya University, Frocho, Chikusaku, Nagoya, 464-8602, Japan
E-mail address: `yuya.mizuno@math.nagoya-u.ac.jp`